

HDAC4 Monoclonal antibody

Catalog Number: 66838-1-Ig 7 Publications

Basic Information

Catalog Number:	66838-1-Ig	GenBank Accession Number:	BC039904	Purification Method:	Protein A purification
Size:	1500 µg/ml	GeneID (NCBI):	9759	CloneNo.:	1D2G12
Source:	Mouse	UNIPROT ID:	P56524	Recommended Dilutions:	WB 1:1000-1:3000 IHC 1:300-1:1200
Isotype:	IgG2a	Full Name:	histone deacetylase 4		
Immunogen Catalog Number:	AG11345	Calculated MW:	972 aa, 106 kDa		
		Observed MW:	140 kDa		

Applications

Tested Applications:	IHC, WB, ELISA	Positive Controls:	
Cited Applications:	WB, IP, IF, ChIP	WB :	HeLa cells, HEK-293 cells, MCF-7 cells, Jurkat cells
Species Specificity:	Human	IHC :	human breast cancer tissue,
Cited Species:	human, mouse		
Note-IHC: suggested antigen retrieval with TE buffer pH 9.0; (*) Alternatively, antigen retrieval may be performed with citrate buffer pH 6.0			

Background Information

HDAC4, also named as HDACA, belongs to the histone deacetylase family. HDAC4 is responsible for the deacetylation of lysine residues on the N-terminal part of the core histones (H2A, H2B, H3 and H4). Histone deacetylation gives a tag for epigenetic repression and plays an important role in transcriptional regulation, cell cycle progression and developmental events. Histone deacetylases act via the formation of large multiprotein complexes. HDAC4 is involved in muscle maturation via its interaction with the myocyte enhancer factors such as MEF2A, MEF2C and MEF2D.

Notable Publications

Author	Pubmed ID	Journal	Application
Yongyun Li	34536440	J Mol Cell Cardiol	IF
Wei Zhan	32404892	Oncogenesis	WB
Wei Jie	35132916	Virulence	IF

Storage

Storage:
Store at -20°C. Stable for one year after shipment.
Storage Buffer:
PBS with 0.02% sodium azide and 50% glycerol pH 7.3.
Aliquoting is unnecessary for -20°C storage

For technical support and original validation data for this product please contact:


T: 4006900926

E: Proteintech-CN@ptglab.com


W: ptgcn.com

This product is exclusively available under Proteintech Group brand and is not available to purchase from any other manufacturer.

Selected Validation Data



Various lysates were subjected to SDS PAGE followed by western blot with 66838-1-Ig (HDAC4 antibody) at dilution of 1:3000 incubated at room temperature for 1.5 hours.



Immunohistochemical analysis of paraffin-embedded human breast cancer tissue slide using 66838-1-Ig (HDAC4 antibody) at dilution of 1:600 (under 40x lens). Heat mediated antigen retrieval with Tris-EDTA buffer (pH 9.0).