For Research Use Only

EPO Monoclonal antibody

Catalog Number:66975-1-lg 4 Publications



Basic Information

Catalog Number:

GenBank Accession Number: BC093628

Purification Method: Protein A purification

66975-1-lg Size:

GeneID (NCBI):

CloneNo.:

1500 µg/ml Source:

2056 UNIPROT ID:

1A11F3

Mouse Isotype: IgG2a

P01588 Full Name: erythropoietin Recommended Dilutions: WB 1:5000-1:50000 IHC 1:50-1:500

Immunogen Catalog Number:

Calculated MW:

AG23054 193 aa, 21 kDa

Applications

Tested Applications: IHC, WB, ELISA

Cited Applications:

WB

WB: recombinant human EPO (cat.NO. HZ-1168), IHC: human liver tissue, mouse kidney tissue

Positive Controls:

Species Specificity: Human, mouse **Cited Species:**

human, rat, mouse

Note-IHC: suggested antigen retrieval with TE buffer pH 9.0; (*) Alternatively, antigen retrieval may be performed with citrate buffer pH 6.0

Background Information

Erythropoietin (Epo) is a member of the EPO/TPO family and encodes a secreted, glycosylated cytokine composed of four alpha helical bundles. The protein is found in the plasma and regulates red cell production by promoting erythroid differentiation and initiating hemoglobin synthesis. Its effect is realized by binding erythropoietin receptor (EpoR) expressed on erythroid progenitor cells. EpoR, is a glycoprotein expressed on megakaryocytes, erythroid progenitors and endothelial cells. Epo also has neuroprotective activity against a variety of potential brain injuries and antiapoptotic functions in several tissue types.

Notable Publications

Author	Pubmed ID	Journal	Application
Takeshi Ike	36074918	Am J Physiol Renal Physiol	WB
Meng Wang	34268128	Front Oncol	WB
Hari Prasad Osuru	35928940	Front Cardiovasc Med	WB

Storage


Storage:

Store at -20°C. Stable for one year after shipment.

PBS with 0.02% sodium azide and 50% glycerol pH 7.3.

Aliquoting is unnecessary for -20°C storage


Selected Validation Data



50 ng recombinant human EPO (cat.NO. HZ-1168) was subjected to SDS-PAGE under reducing condition followed by western blot with 66975-1-lg (EPO antibody) at dilution of 1:10000.



Immunohistochemical analysis of paraffinembedded human liver tissue slide using 66975-1-Ig (EPO antibody) at dilution of 1:200 (under 10x lens. Heat mediated antigen retrieval with Tris-EDTA buffer (pH 9.0).



Immunohistochemical analysis of paraffinembedded human liver tissue slide using 66975-1-Ig (EPO antibody) at dilution of 1:200 (under 40x lens. Heat mediated antigen retrieval with Tris-EDTA buffer (pH 9.0).