

HOXA7 Monoclonal antibody

Catalog Number: 67112-1-Ig 3 Publications

Basic Information

Catalog Number:

67112-1-Ig

Size:

2500 µg/ml

Source:

Mouse

Isotype:

IgG1

Immunogen Catalog Number:

AG19025

GenBank Accession Number:

BC148692

GeneID (NCBI):

3204

UNIPROT ID:

P31268

Full Name:

homeobox A7

Calculated MW:

230 aa, 25 kDa

Observed MW:

20 kDa

Purification Method:

Protein G purification

CloneNo.:

2A7H6

Recommended Dilutions:

WB 1:1000-1:6000

IF 1:50-1:500

Applications

Tested Applications:

IF/ICC, WB, ELISA

Cited Applications:

WB, IF, IHC

Species Specificity:

Human

Cited Species:

human, mouse

Positive Controls:

WB : THP-1 cells, HepG2 cells, SMMC-7721 cells, HT-29 cells, HL-60 cells, K-562 cells

IF : HepG2 cells,

Background Information

HOX genes play a fundamental role in the development of the vertebrate central nervous system, heart, axial skeleton, limbs, gut, urogenital tract and external genitalia. The homeobox gene Hoxa-1 is transcriptionally regulated by retinoic acid (RA) and encodes a transcription factor, which has been shown to play important roles in cell differentiation and embryogenesis. Hoxa-1 is also expressed in cancers, such as mammary tumors, though it is not expressed in normal gland or in precancerous mammary tissues. At embryonic stages, Hoxa-2 is expressed in the mesenchyme and epithelial cells of palate, however its expression is restricted to the tips of the growing palatal shelves. Hoxa-2 protein is predominantly expressed in the nuclei of cells in the ventral mantle region of the developing embryo. In the developing and adult mouse spinal cord, Hoxa-2 protein may contribute to dorsal-ventral patterning and/or to the specification of neuronal phenotype. Hoxa-7 functions as a potent transcriptional repressor and its action as such requires several domains, including both activator and repressor regions. Hoxa-7 is expressed in the fetal liver, lung, skeletal muscle, kidney, pancreas and placenta.

Notable Publications

Author	Pubmed ID	Journal	Application
Jun-Pu Zha	32650806	J Orthop Surg Res	WB
Jun Li	34978042	Reprod Sci	WB
Anqi Feng	37248653	Cancer Sci	WB, IHC, IF

Storage

Storage:

Store at -20°C.

Storage Buffer:

PBS with 0.02% sodium azide and 50% glycerol pH 7.3.

Aliquoting is unnecessary for -20°C storage


For technical support and original validation data for this product please contact:

T: 4006900926


E: Proteintech-CN@ptglab.comW: ptgcn.com

This product is exclusively available under Proteintech Group brand and is not available to purchase from any other manufacturer.

Selected Validation Data



THP-1 cells were subjected to SDS PAGE followed by western blot with 67112-1-Ig (HOXA7 antibody) at dilution of 1:3000 incubated at room temperature for 1.5 hours.



Immunofluorescent analysis of (4% PFA) fixed HepG2 cells using 67112-1-Ig (HOXA7 antibody) at dilution of 1:100 and CoraLite488-Conjugated AffiniPure Goat Anti-Mouse IgG(H+L).