For Research Use Only

SOX9 Monoclonal antibody

Catalog Number:67439-1-lg Featured Product

43 Publications

BC007951

6662

GeneID (NCBI):

GenBank Accession Number:



Basic Information

Catalog Number: 67439-1-lg

Size: 2000 μg/ml Source:

UNIPROT ID: Mouse P48436 Full Name: Isotype:

lgG1 SRY (sex determining region Y)-box 9 Calculated MW:

Immunogen Catalog Number: AG28620 509 aa, 56 kDa Observed MW: 56 kDa

Applications

Tested Applications: FC, WB, ELISA Cited Applications: WB, IF, FC, IHC Species Specificity:

Human, mouse **Cited Species:**

human, rat, mouse, rabbit, pig, canine

Positive Controls:

WB: HT-29 cells, Caco-2 cells, human placenta tissue, NCCIT cells

Purification Method:

Protein A purification

Recommended Dilutions:

WB 1:5000-1:50000

CloneNo.:

1H9A6

Background Information

SOX9, also named as Transcription factor SOX-9, is a 509 amino acid protein. Sex-determining region Y (SRY)-box 9 protein (SOX9) is a member of the SOX family of transcription factors (TFs) which are developmental regulators that possess high mobility group (HMG) box DNA binding and transactivation domains. It participates in a variety of functions, such as lineage restriction and terminal diferentiation, through precise temporal and spatial expression patterns that difer between particular cell types and tissues. SOX9 gene has been implicated in different types of cancer as an oncogene; however, it also may behave as a tumor suppressor.

Notable Publications

Author	Pubmed ID	Journal	Application
LinXi Cheng	34489520	Commun Biol	FC
Xiao-Xin Gao	34719207	Front Biosci (Landmark Ed)	WB
Tianwen Ma	36386227	Front Pharmacol	WB

Storage


Storage:

Store at -20°C. Stable for one year after shipment.


PBS with 0.02% sodium azide and 50% glycerol pH 7.3.

Aliquoting is unnecessary for -20°C storage


Selected Validation Data



Various lysates were subjected to SDS PAGE followed by western blot with 67439-1-lg (SOX9 antibody) at dilution of 1:10000 incubated at room temperature for 1.5 hours. The membrane was stripped and reblotted with HRP-conjugated GAPDH Monoclonal antibody (HRP-60004) as loading control



human placenta tissue were subjected to SDS PAGE followed by western blot with 67439-1-lg (SOX9 antibody) at dilution of 1:3000 incubated at room temperature for 1.5 hours.



1X10^6 HeLa cells were intracellularly stained with 0.4 ug Anti-Human SOX9 (67439-1-Ig, Clone:1H9A6) and CoraLite® 488-Conjugated Affini Pure Goat Anti-Mouse IgG(H+L) at dilution 1:1000 (red), or 0.4 ug Mouse IgG1 Isotype Control (MOPC-21) (65124-1-Ig, Clone: MOPC-21) (blue). Cells were fixed and permeabilized with Transcription Factor Staining Buffer Kit (PF00011).