

For Research Use Only

CCT4 Monoclonal antibody

Catalog Number: 67455-1-Ig



Basic Information

Catalog Number:

67455-1-Ig

Size:

1000 µg/ml

Source:

Mouse

Isotype:

IgG1

Immunogen Catalog Number:

AG17084

GenBank Accession Number:

BC106934

GeneID (NCBI):

10575

UNIPROT ID:

P50991

Full Name:

chaperonin containing TCP1, subunit

4 (delta)

Calculated MW:

539 aa, 58 kDa

Observed MW:

55-58 kDa

Purification Method:

Protein G purification

CloneNo.:

3C6A6

Recommended Dilutions:

WB 1:5000-1:50000

IF 1:200-1:800

Applications

Tested Applications:

WB, IF/ICC, ELISA

Species Specificity:

Human, Mouse, Rabbit, Pig

Positive Controls:

WB : LNCaP cells, HeLa cells, HEK-293 cells, HepG2 cells, Jurkat cells, pig brain tissue, rabbit brain tissue, mouse brain tissue

IF : U2OS cells,

Background Information

T-complex protein 1 subunit delta (CCT4) is a component of the chaperonin-containing T-complex (TRiC), a molecular chaperone complex that assists the folding of proteins upon ATP hydrolysis (PMID:25467444). As part of the TRiC complex, it may play a role in regulating telomere maintenance (PMID:25467444) and assembly of BBSome, a complex involved in ciliogenesis regulating transports vesicles to the cilia (PMID:20080638). CCT4 has 2 isoforms produced by alternative splicing with the MW of 55 kDa and 58 kDa.

Storage

Storage:

Store at -20°C. Stable for one year after shipment.

Storage Buffer:

PBS with 0.1% sodium azide and 50% glycerol pH 7.3.

Aliquoting is unnecessary for -20°C storage

For technical support and original validation data for this product please contact:


T: 4006900926

E: Proteintech-CN@ptglab.com

W: ptgcn.com

This product is exclusively available under Proteintech Group brand and is not available to purchase from any other manufacturer.

Selected Validation Data



Immunofluorescent analysis of (-20°C Methanol) fixed U2OS cells using CCT4 antibody (67455-1-Ig, Clone: 3C6A6) at dilution of 1:400 and CoraLite®488-Conjugated AffiniPure Goat Anti-Mouse IgG(H+L).

Various lysates were subjected to SDS PAGE followed by western blot with 67455-1-Ig (CCT4 antibody) at dilution of 1:20000 incubated at room temperature for 1.5 hours. The membrane was stripped and reblotted with HRP-conjugated GAPDH Monoclonal antibody (HRP-60004) as loading control.