

For Research Use Only

NeutraKine® IL-3 Mouse McAb

Catalog Number: 69004-1-Ig



Basic Information

Catalog Number:	69004-1-Ig	GenBank Accession Number:	Purification Method:
Concentration:		GeneID (NCBI):	Protein G purification
Source:	Mouse	3562	CloneNo.:
Isotype:	IgG1	Full Name:	3C11G7
Immunogen Catalog Number:	HZ-1074	interleukin 3 (colony-stimulating factor, multiple)	

Applications

Tested Applications:	Positive Controls:
Neutralization, ELISA	Neutralization : TF-1 cells,
Species Specificity:	
human	

Background Information

Interleukin-3 (IL-3) is a multilineage hematopoietic growth factor that promotes the proliferation, differentiation and survival of early multilineage hematopoietic progenitors. In particular, this cytokine plays a key role in stimulating the proliferation and survival of myeloid precursors. It is involved in a variety of cell activities such as cell growth, differentiation and apoptosis. This cytokine has been shown to also possess neurotrophic activity, and it may be associated with neurologic disorders (PMID: 24103757;2698876;2809196).

This antibody can be used to neutralize the bioactivity of IL-3.

Storage

Storage:
Lyophilized antibodies are stable for 1 year from the date of receipt if stored between (-20°C) and (-80°C). Upon reconstitution we recommend that the solution can be stored at (4°C) for short term or at (-20°C) to (-80°C) for long term. Repeated freeze thaw cycles should be avoided with reconstituted products.

Storage Buffer:
Sterile PBS, pH7.4

Aliquoting is unnecessary for -20°C storage

For technical support and original validation data for this product please contact:


T: 4006900926

E: Proteintech-CN@ptglab.com


W: ptgcn.com

This product is exclusively available under Proteintech Group brand and is not available to purchase from any other manufacturer.

Selected Validation Data



Recombinant human IL-3 (Cat.NO. HZ-1074) stimulates proliferation of TF-1 cell line (human erythroleukemic cell line) in a dose-dependent manner (blue curve, refer to bottom X-left Y). The activity of human IL-3 (10 ng/mL HZ-1074) is neutralized by mouse anti-human IL-3 monoclonal antibody 69004-1-Ig at serial dose (red curve, refer to top X-right Y). The ND50 is typically 40-70 ng/mL.



Recombinant human IL-3 (Cat.NO. HZ-1074) stimulates proliferation of TF-1 cell line (human erythroleukemic cell line) in a dose-dependent manner (blue curve, refer to bottom X-left Y). The activity of human IL-3 (0.5 ng/mL HZ-1074) is neutralized by mouse anti-human IL-3 monoclonal antibody 69004-1-Ig at serial dose (red curve, refer to top X-right Y). The ND50 is typically 0.8-2.4 ng/mL.