### For Research Use Only

# Cytokeratin 18 Recombinant antibody

proteintech®

Antibodies | ELISA kits | Proteins

www.ptglab.com

Catalog Number:82817-5-RR

**Basic Information** 

**Applications** 

Catalog Number: GenBank Accession Number:

82817-5-RR BC000180 Size: GeneID (NCBI): 500 μg/ml 3875

Source: UNIPROT ID:
Rabbit P05783
Isotype: Full Name:
IgG keratin 18

AG1260 48 kDa

Immunogen Catalog Number:

Observed MW: 48 kDa

Tested Applications:

Calculated MW:

IHC, WB, ELISA

Species Specificity:
Human, mouse, rat

Note-IHC: suggested antigen retrieval with TE buffer pH 9.0; (\*) Alternatively, antigen retrieval may be performed with citrate buffer pH 6.0 WB: A431 cells, A549 cells, HCT 116 cells, HepG2 cells, K-562 cells, mouse liver tissue, rat liver tissue

Positive Controls:

**Purification Method:** 

CloneNo.:

IHC 1:50-1:500

Protein A purification

Recommended Dilutions: WB 1:2000-1:10000

IHC: human appendicitis tissue, human liver tissue, human renal cell carcinoma tissue, human breast

# **Background Information**

Cytokeratin 18, also known as CK18, is a type I intermediate filament protein which is found primarily in non squamous epithelia and is present in a majority of adenocarcinomas and ductal carcinomas but not in squamous cell carcinomas. CK18 and its coexpressed complementary type II keratin partner, CK8, are persistently expressed in a variety of adult epithelial organs, such as the liver, lung, kidney, pancreas, gastrointestinal tract, and mammary gland, and are also expressed by cancers that arise from these tissues. In the absence of CK8, the CK18 protein is degraded and keratin intermediate filaments are not formed.

#### Storage

Storage


Store at -20°C. Stable for one year after shipment.

Storage Buffe


PBS with 0.02% sodium azide and 50% glycerol pH 7.3.

Aliquoting is unnecessary for -20°C storage


## Selected Validation Data



Various lysates were subjected to SDS PAGE followed by western blot with 82817-5-RR (KRT 18 antibody) at dilution of 1:5000 incubated at room temperature for 1.5 hours.



Immunohistochemical analysis of paraffinembedded human appendicitis tissue slide using 82817-5-RR (Cytokeratin 18 antibody) at dilution of 1:200 (under 10x lens). Heat mediated antigen retrieval with Tris-EDTA buffer (pH 9.0).



Immunohistochemical analysis of paraffinembedded human appendicitis tissue slide using 82817-5-RR (Cytokeratin 18 antibody) at dilution of 1:200 (under 40x lens). Heat mediated antigen retrieval with Tris-EDTA buffer (pH 9.0).