For Research Use Only

Cytokeratin 13 Recombinant antibody

Size:

800 μ g/ml

proteintech®
Antibodies | ELISA kits | Proteins
www.ptglab.com

Catalog Number:83058-1-RR

Basic Information

Catalog Number: GenBank Accession Number: 83058-1-RR BC002661

BC002661 GeneID (NCBI): 3860

Source: UNIPROT ID:
Rabbit P13646

Isotype: Full Name:
IgG keratin 13

Immunogen Catalog Number: Calculated MW:

AG0217 50 kDa

721/ 30 KD

Observed MW: 50 kDa

Purification Method:

Protein A purification

CloneNo.: 230366B10

Recommended Dilutions: WB 1:2000-1:16000 IHC 1:500-1:2000 IF 1:125-1:500

Applications

Tested Applications: WB, IF/ICC, IHC, ELISA Species Specificity:

Human, mouse, rat

Note-IHC: suggested antigen retrieval with TE buffer pH 9.0; (*) Alternatively, antigen retrieval may be performed with citrate buffer pH 6.0 Positive Controls:

WB: HaCaT cells, mouse skin tissue, rat skin tissue, rat thymus tissue

IHC: human urothelial carcinoma tissue, rat skin tissue, mouse skin tissue, human oesophagus cancer tissue, human cervical cancer tissue

IF: HaCaT cells,

Background Information

Keratin 13 is a member of the keratin family. The keratins are intermediate filament proteins responsible for the structural integrity of epithelial cells and are subdivided into cytokeratins and hair keratins. Most of the type I cytokeratins consist of acidic proteins which are arranged in pairs of heterotypic keratin chains. This type I cytokeratin is paired with keratin 4 and expressed in the suprabasal layers of non-cornified stratified epithelia. Mutations in keratin 13 gene and keratin 4 have been associated with the autosomal dominant disorder White Sponge Nevus. The type I cytokeratins are clustered in a region of chromosome 17q21.2.

Storage


Storage

Store at -20°C. Stable for one year after shipment. Storage Buffer:


PBS with 0.02% sodium azide and 50% glycerol pH 7.3.

Aliquoting is unnecessary for -20° C storage


Selected Validation Data




Various lysates were subjected to SDS PAGE followed by western blot with 83058-1-RR (KRT 13 antibody) at dilution of 1:8000 incubated at room temperature for 1.5 hours.



Immunohistochemical analysis of paraffinembedded human urothelial carcinoma tissue slide using 83058-1-RR (Cytokeratin 13 antibody) at dilution of 1:1000 (under 10x lens). Heat mediated antigen retrieval with Tris-EDTA buffer (pH 9.0).



Immunohistochemical analysis of paraffinembedded human urothelial carcinoma tissue slide using 83058-1-RR (Cytokeratin 13 antibody) at dilution of 1:1000 (under 40x lens). Heat mediated antigen retrieval with Tris-EDTA buffer (pH 9.0).



Immunofluorescent analysis of (-20°C Methanol) fixed HaCaT cells using Cytokeratin 13 antibody (83058-1-RR, Clone: 230366B10) at dilution of 1:250 and CoraLite®488-Conjugated AffiniPure Goat Anti-Rabbit IgG(H+L) (SA00013-2).