For Research Use Only

CoraLite® Plus 488-conjugated STAT3 Polyclonal antibody



Catalog Number: CL488-10253

Featured Product

Basic Information

Catalog Number: CL488-10253

1000 µg/ml Source: Rabbit Isotype:

Immunogen Catalog Number:

AG0360

GenBank Accession Number:

BC000627 GeneID (NCBI): 6774 UNIPROT ID: P40763 Full Name:

signal transducer and activator of transcription 3 (acute-phase response

factor)

Calculated MW: 770 aa, 88 kDa Observed MW: 88 kDa Purification Method:

Antigen affinity purification Excitation/Emission maxima

wavelengths: 493 nm / 522 nm

Applications

Tested Applications:

FC (Intra)

Species Specificity: human, mouse, rat

Background Information

Signal transducer and activator of transcription 3 (acute-phase response factor) (STAT3, synonyms: APRF, FLJ20882, MGC16063) is a member of the STAT protein family. In response to cytokines and growth factors, STAT family members are phosphorylated by the receptor associated kinases, and then form homo- or heterodimers that translocate to the cell nucleus where they act as transcription activators. STAT3 is activated through phosphorylation in response to various cytokines and growth factors including IFNs, EGF, IL5, IL6, HGF, LIF and BMP2. STAT3 mediates the expression of a variety of genes in response to cell stimuli, and thus plays a key role in many cellular processes such as cell growth and apoptosis. The small GTPase Rac1 has been shown to bind and regulate the activity of STAT3. This antibody is a rabbit polyclonal antibody raised against residues near the N terminus of human STAT3. STAT3 exists three isoforms and the molecular weight of each isoform respectively is 83 kDa, 87 kDa and 88 kDa.

Storage

Storage:


Store at -20°C. Avoid exposure to light.

Storage Buffe

PBS with 50% Glycerol, 0.05% Proclin300, 0.5% BSA, pH 7.3.

Aliquoting is unnecessary for -20°C storage

Selected Validation Data



1X10^6 HeLa cells were intracellularly stained with 0.4 ug CoraLite® Plus 488 Anti-Human STAT3 (CL488-10253) (red), or 0.4 ug Control Antibody. Cells were fixed and permeabilized with Transcription Factor Staining Buffer Kit (PF00011).