For Research Use Only

CoraLite® Plus 488-conjugated c-MYC Polyclonal antibody



Catalog Number: CL488-10828

Featured Product

Basic Information

Catalog Number: CL488-10828

1000 µ g/ml
Source:
Rabbit
Isotype:

Immunogen Catalog Number:

AG1263

bbit

oncogene homolog (avian)

Calculated MW:

BC000141

4609

P01106

GeneID (NCBI):

UNIPROT ID:

Full Name:

GenBank Accession Number:

v-myc myelocytomatosis viral

49 kDa

Observed MW:
62-65 kDa, 50 kDa

Purification Method:

Antigen affinity purification

Recommended Dilutions: IF 1:300-1:1200

Excitation/Emission maxima

wavelengths: 493 nm / 522 nm

Applications

Tested Applications: FC (Intra), IF/ICC Species Specificity:

Species Specificity human, mouse Positive Controls:

IF: HEK-293 cells,

Background Information

MYC contains one basic helix-loop-helix (bHLH) domain. This protein is a multifunctional, nuclear phosphoprotein that plays a role in cell cycle progression, apoptosis and cellular transformation. It functions as a transcription factor that regulates transcription of specific target genes. It seems to activate the transcription of growth-related genes. It binds DNA in a non-specific manner, yet also specifically recognizes the core sequence 5'-CAC[GA]TG-3'. Mutations, overexpression, rearrangement and translocation of this gene have been associated with a variety of hematopoietic tumors, leukemias and lymphomas, including Burkitt lymphoma. This antibody is a rabbit polyclonal antibody raised against recombinant protein of human MYC.The 50kDa band recognized by antibody is the native form of MYC, while the other bands, between 60-70kDa, are the phosphorylated form of MYC [PMID: 12189186]

Storage

Storage:

Store at -20°C. Avoid exposure to light.

Storage Buffer


PBS with 50% Glycerol, 0.05% Proclin300, 0.5% BSA, pH 7.3.

Aliquoting is unnecessary for -20°C storage

Selected Validation Data



Immunofluorescent analysis of (4% PFA) fixed HEK-293 cells using CoraLite® Plus 488 c-MYC antibody (CL488-10828) at dilution of 1:600.



1X10^6 HeLa cells were intracellularly stained with 0.4 ug CoraLite® Plus 488 Anti-Human c-MYC (CL488-10828) (red), or 0.4 ug Control Antibody. Cells were fixed and permeabilized with Transcription Factor Staining Buffer Kit (PF00011).