For Research Use Only

## CoraLite® Plus 488-conjugated NR3C2 Polyclonal antibody



**Purification Method:** 

IF 1:50-1:500

wavelengths: 493 nm / 522 nm

Antigen affinity purification

Excitation/Emission maxima

Recommended Dilutions:

Catalog Number: CL488-21854

**Featured Product** 

**Basic Information** 

Catalog Number: CL488-21854

1000 µg/ml Source: Rabbit

Isotype:

Immunogen Catalog Number:

AG16410

**Tested Applications:** Species Specificity:

human, mouse

GenBank Accession Number:

BC111758 GeneID (NCBI): 4306

**UNIPROT ID:** P08235 Full Name:

member 2

94-110 kDa

nuclear receptor subfamily 3, group C,

Calculated MW: 984 aa, 107 kDa Observed MW:

Positive Controls:

IF: HepG2 cells,

## **Background Information**

Nuclear receptor subfamily 3 group C member 2 (NR3C2), also known as Mineralocorticoid receptor (MCR or MR), is a member of the steroid/thyroid/retinoic nuclear hormone receptor superfamily that has been shown to activate gene transcription in response to aldosterone binding. Regulation of the mineral ocorticoid receptors occurs through either receptor down-regulation (negative autoregulation) or hormone-mediated upregulation (positive autoregulation). MCR association with HSP 90 appears to be required for hormone binding to MCR and subsequent MCR activation.

Storage


**Applications** 

Store at -20°C. Avoid exposure to light. Stable for one year after shipment.

PBS with 50% Glycerol, 0.05% Proclin300, 0.5% BSA, pH 7.3.

Aliquoting is unnecessary for -20°C storage

## Selected Validation Data



Immunofluorescent analysis of (4% PFA) fixed HepG2 cells using CoraLite® Plus 488 NR3C2 antibody (CL488-21854) at dilution of 1:200.