For Research Use Only

CoraLite® Plus 488-conjugated STAT3 Monoclonal antibody



Featured Product


Antibodies | ELISA kits | Proteins www.ptglab.com

Basic Information	Catalog Number: CL488-60199	GenBank Accession Number: BC000627	Purification Method: Protein A purification
	Size: 1000 µg/ml	GeneID (NCBI): 6774	CloneNo.: 3G2D12
	Source: Mouse	UNIPROT ID: P40763	Recommended Dilutions: IF 1:50-1:500
	Isotype: IgG2a Immunogen Catalog Number: AG0360	Full Name:Excitation/Emission maximasignal transducer and activator ofwavelengths:transcription 3 (acute-phase response 493 nm / 522 nmfactor)Calculated MW:770 aa, 88 kDa	
		Applications	Tested Applications:
FC (Intra), IF/ICC Species Specificity: human, mouse, rat	IF : LO2 cells, HepG2 cells		
Background Information	Signal transducer and activator of transcription 3 (acute-phase response factor) (STAT3, synonyms: APRF, FLJ20882, MGC16063) is a member of the STAT protein family. In response to cytokines and growth factors, STAT family members are phosphorylated by the receptor associated kinases, and then form homo- or heterodimers that translocate to the cell nucleus where they act as transcription activators. STAT3 is activated through phosphorylation in response to various cytokines and growth factors including IFNs, EGF, IL5, IL6, HGF, LIF and BMP2. STAT3 mediates the expression of a variety of genes in response to cell stimuli, and thus plays a key role in many cellular processes such as cell growth and apoptosis. The small GTPase Rac1 has been shown to bind and regulate the activity of STAT3. This antibody is a mouse monoclonal antibody raised against residues near the N terminus of human STAT3. STAT3 exists three isoforms and the molecular weight of each isoform respectively is 83 kDa, 87 kDa and 88 kDa.		
Storage	Storage: Store at -20°C. Avoid exposure to light. Stable for one year after shipment. Storage Buffer: PBS with 50% Glycerol, 0.05% Proclin300, 0.5% BSA, pH 7.3. Aliquoting is unnecessary for -20°C storage		


For technical support and original validation data for this product please contact:T: 4006900926E: Proteintech-CN@ptglab.comW: ptgcn.com

This product is exclusively available under Proteintech Group brand and is not available to purchase from any other manufacturer.

Selected Validation Data



Immunofluorescent analysis of (-20°C Ethanol) fixed L02 cells using CoraLite® Plus 488conjugated STAT3 antibody (CL488-60199, Clone: 3G2D12) at dilution of 1:100, CL594-Phalloidin (red).



1X10^6 HepG2 cells were intracellularly stained with 0.4 ug CoraLite® Plus 488 Anti-Human STAT3 (CL488-60199, Clone:3G2D12) (red), or 0.4 ug Mouse 1gG2a 1sotype Control (CL488-66360-2, Clone: K11A1B2A2) (blue). Cells were fixed with 4% PFA and permeabilized with Flow Cytometry Perm Buffer (PF00011-C).