For Research Use Only

CoraLite®488-conjugated Vimentin Monoclonal antibody



Purification Method:

Protein G purification

Recommended Dilutions:

Excitation/Emission maxima

CloneNo.:

IF 1:200-1:800

wavelengths: 491 nm / 516 nm

3H9D1

Catalog Number: CL488-60330

1 Publications

Basic Information

Applications

Catalog Number:

CL488-60330

1000 µg/ml

Source:

Mouse Isotype: lgG1

Immunogen Catalog Number:

AG0489

Calculated MW:

UNIPROT ID:

Full Name:

vimentin

P08670

BC000163

7431

GeneID (NCBI):

ENSEMBL Gene ID:

ENSG00000026025

466 aa, 54 kDa

GenBank Accession Number:

Tested Applications:

FC (Intra), IF/ICC

Cited Applications:

IF, IHC

mouse

Species Specificity: human, mouse, rat Cited Species:

Positive Controls:

IF: HepG2 cells, human tonsillitis tissue

Background Information

Vimentin, also named as VIM, belongs to the intermediate filament family. Vimentin is class-III intermediate filaments found in various non-epithelial cells, especially mesenchymal cells. Vimentin is important for stabilizing the architecture of the cytoplasm. Monocyte-derived macrophages secrete vimentin into the extracellular space in vitro. Secretion of vimentin was enhanced by the proinflammatory cytokine tumor necrosis factor-alpha (TNFA; 191160) and inhibited by the antiinflammatory cytokine IL10 (124092), suggesting that vimentin is involved in the immune response. Vimentin has specialized functions that contribute to specific dynamic cellular processes. As a phosphoprotein, 55-60 kDa of vimentin proteins can be observed due to the different phosphorylation level. The antibody is conjugated with CL488, Ex/Em 488 nm/515 nm.

Notable Publications

Author	Pubmed ID	Journal	Application
Feng Zheng	33760044	J Mol Cell Biol	IHC, IF


Storage

Store at -20°C. Avoid exposure to light. Stable for one year after shipment.


PBS with 50% Glycerol, 0.05% Proclin300, 0.5% BSA, pH 7.3.

Aliquoting is unnecessary for -20°C storage

Selected Validation Data



Immunofluorescent analysis of (-20°C Methanol) fixed HepG2 cells using CoraLite@488 Vimentin antibody (CL488-60330, Clone: 3H9D1) at dilution of 1:400.



1X10^6 Jurkat cells were intracellularly stained with 0.4 ug Coralite® 488 Anti-Human Vimentin (CL488-60330, Clone:3H9D1) (red), or 0.4 ug Mouse IgG1 Isotype Control (CL488-66360, Clone: T1F8D3F10) (blue). Cells were fixed with 4% PFA and permeabilized with Flow Cytometry Perm Buffer (PF00011-C).



Immunofluorescent analysis of (-20°C Methanol) fixed HepG2 cells using CoraLite@488 Vimentin antibody (CL488-60330, Clone: 3H9D1) at dilution of 1:400.