## For Research Use Only

## CoraLite® Plus 488-conjugated CD8a Monoclonal antibody

Catalog Number:CL488-66868

1 Publications




| Basic Information      | Catalog Number:<br>CL488-66868   | GenBank Accession Number:<br>BC025715   | Purification Method:<br>Protein G purification   |
|------------------------|--|---|--|
|                        | Size:  | GenelD (NCBI):  | CloneNo.:  |
|                        | 1000 µg/ml   | 925   | 1G2B10   |
|                        | Source:<br>Mouse   | ENSEMBL Gene ID:<br>ENSG00000153563   | Recommended Dilutions:<br>IF 1:50-1:500  |
|                        | Isotype:<br>IgG1<br>Immunogen Catalog Number:<br>AG11111   | UNIPROT ID:<br>P01732   | Excitation/Emission maxima<br>wavelengths:<br>493 nm / 522 nm  |
|                        |  | Full Name:<br>CD8a molecule   |  |
|                        |  | Calculated MW:<br>235 aa, 26 kDa  |  |
| Applications           | Tested Applications:<br>IF/ICC,IF-P  | Positive Controls:<br>IF : human tonsillitis tissue,  |  |
|                        | Cited Applications:<br>IHC   |   |  |
|                        | Species Specificity:   |   |  |
|                        | Human  |   |  |
| Background Information | CD8 is a transmembrane glycopro<br>also be found on natural killer cel<br>cell receptor (TCR). Both CD8 and<br>context of class I MHC molecules.   | ls, cortical thymocytes, and dendritic of TCR recognize antigens displayed by a CD8 plays a role in T cell development  |  |
|                        | CD8 is a transmembrane glycopro<br>also be found on natural killer cel<br>cell receptor (TCR). Both CD8 and<br>context of class I MHC molecules.<br>antibody is raised against 23-183<br>CL488, Ex/Em 488 nm/515 nm. | ls, cortical thymocytes, and dendritic of TCR recognize antigens displayed by a CD8 plays a role in T cell development  | cells. CD8 serves as a co-receptor for the<br>an antigen presenting cell (APC) in the<br>nt and activation of mature T cells. This |
| Background Information | CD8 is a transmembrane glycopro<br>also be found on natural killer cel<br>cell receptor (TCR). Both CD8 and<br>context of class I MHC molecules.<br>antibody is raised against 23-183<br>CL488, Ex/Em 488 nm/515 nm. | ls, cortical thymocytes, and dendritic of<br>TCR recognize antigens displayed by a<br>CD8 plays a role in T cell developmer<br>aa of human CD8a, the alpha chain of | nt and activation of mature T cells. This<br>CD8 molecule, and is conjugated with  |

For technical support and original validation data for this product please contact:T: 4006900926E: Proteintech-CN@ptglab.comW: ptgcn.com

This product is exclusively available under Proteintech Group brand and is not available to purchase from any other manufacturer.

## Selected Validation Data



Immunofluorescent analysis of (4% PFA) fixed human tonsillitis tissue using CoraLite® Plus 488 CD8a antibody (CL488-66868, Clone: 1G2B10) at dilution of 1:200, CoraLite® 594 CD11c/Integrin Alpha X antibody (CL594-60258, Clone: 2F1C10, red).