For Research Use Only

CoraLite®555-conjugated Galectin-3 Monoclonal antibody


Catalog Number: CL555-60207

Featured Product


Basic Information	Catalog Number: CL555-60207	GenBank Accession Number: BC001120	Purification Method: Protein A purification
	Size: 1000 μg/ml	GeneID (NCBI): 3958	CloneNo.: 1C1B2
	Source: Mouse	UNIPROT ID: P17931	Recommended Dilutions: IF 1:50-1:500
	lsotype: lgG2b Immunogen Catalog Number: AG6753	Full Name:Excitation/Emission maximalectin, galactoside-binding, soluble, 3 wavelengths:Calculated MW:26 kDa	
			557 nm / 570 nm
		Observed MW: 31 kDa	
Applications	Tested Applications:	Positive Controls: IF : A431 cells,	
	FC, IF/ICC Species Specificity: human, mouse, rat		
Background Information	Galectins are a family of animal lectins defined by shared characteristic amino-acid sequences and affinity for β - galactose-containing oligosac-charides (PMID: 8063692). Galectin-3, a member of the β -galactoside-binding proteins, contains one carbohydrate recognition domain (CRD) and a proline- and glycine-rich N-terminal domain through which is able to form oligomers (PMID: 14758078). Galectin-3 is widely expressed in many normal tissues and a variety of tumors. It is found intracellularly in nucleus and cytoplasm or secreted outside of cell, being present on the cell surface or in the extracellular space (PMID: 16478649). Galectin-3 is involved in various biological processes including cell growth, adhesion, differentiation, apoptosis, angiogenesis, immune response, neoplastic transformation and metastasis (PMID: 16478649; 14758078).		
Storage	Storage: Store at -20°C. Avoid exposure to Storage Buffer: PBS with 50% Glycerol, 0.05% Pro Aliquoting is unnecessary for -20°		ent.

This product is exclusively available under Proteintech Group brand and is not available to purchase from any other manufacturer.

Selected Validation Data



Immunofluorescent analysis of (-20°C Methanol) fixed A431 cells using CoraLite®555 Galectin-3 antibody (CL555-60207, Clone: 1C1B2) at dilution of 1:200.



1X10⁶ HeLa cells were intracellularly stained with 0.4 ug CoraLite®555 Anti-Human Galectin-3 (CL555-60207, Clone:1C1B2) (red), or 0.4 ug Control Antibody. Cells were fixed with 4% PFA and permeabilized with Flow Cytometry Perm Buffer (PF00011-C).