

CoraLite®594-conjugated NRF2, NFE2L2 Polyclonal antibody

Catalog Number: CL594-16396

Featured Product

1 Publications

Basic Information

Catalog Number:

CL594-16396

Size:

1000 µg/ml

Source:

Rabbit

Isotype:

IgG

Immunogen Catalog Number:

AG9489

GenBank Accession Number:

BC011558

GeneID (NCBI):

4780

UNIPROT ID:

Q16236

Full Name:

nuclear factor (erythroid-derived 2)-like 2

Calculated MW:

605 aa, 68 kDa

Observed MW:

110 kDa

Purification Method:

Antigen affinity purification

Recommended Dilutions:

IF 1:50-1:500

Excitation/Emission maxima wavelengths:

588 nm / 604 nm

Applications

Tested Applications:

FC (Intra), IF/ICC

Cited Applications:

IF

Species Specificity:

human, mouse, rat

Cited Species:

rat

Positive Controls:

IF : HepG2 cells,

Background Information

NRF2, also named as NFE2L2, belongs to the bZIP family and CNC subfamily. It is a transcription activator that binds to antioxidant response (ARE) elements in the promoter regions of target genes. NRF2 is important for the coordinated up-regulation of genes in response to oxidative stress. It may be involved in the transcriptional activation of genes of the beta-globin cluster by mediating enhancer activity of hypersensitive site 2 of the beta-globin locus control region. Nrf2 is a key player in the regulation of genes encoding for many antioxidative response enzymes. The expression of NRF2 may be induced under oxidative stress (PMID:14567983). In lung cancer, Nrf2 activation in malignant cells has been associated with tumor progression and chemotherapy resistance (PMID:20534738). Identifying patients with abnormal NRF2 expression may be important for selection for chemotherapy in NSCLC. As new investigators break into the emerging field of Nrf2 research, confusion regarding the correct migratory pattern of Nrf2 is causing doubts about the accuracy and reproducibility of published results. This letter provides solid evidence that the actually observed molecular weight of Nrf2 is about 70kDa and 95-110 kDa. (PMID: 22703241).

Notable Publications

Author	Pubmed ID	Journal	Application
Qi Li	36974993	Am J Chin Med	IF

Storage

Storage:

Store at -20°C. Avoid exposure to light.

Storage Buffer:

PBS with 50% Glycerol, 0.05% Proclin300, 0.5% BSA, pH 7.3.

Aliquoting is unnecessary for -20°C storage


For technical support and original validation data for this product please contact:

T: 4006900926


E: Proteintech-CN@ptglab.comW: ptgcn.com

This product is exclusively available under Proteintech Group brand and is not available to purchase from any other manufacturer.


Selected Validation Data




Immunofluorescent analysis of (-20°C Ethanol) fixed HepG2 cells using CoraLite®594 NRF2, NFE2L2 antibody (CL594-16396) at dilution of 1:100.



1X10⁶ MCF-7 cells were intracellularly stained with 0.2 ug CoraLite®594 Anti-Human NRF2, NFE2L2 (CL594-16396) (red), or 0.2 ug Control Antibody. Cells were fixed and permeabilized with Transcription Factor Staining Buffer Kit (PF00011).



Immunofluorescent analysis of (-20°C Methanol) fixed HepG2 cells using CoraLite®594 NRF2, NFE2L2 antibody (CL594-16396) at dilution of 1:200.



1X10⁶ MCF-7 cells were intracellularly stained with 0.4 ug CoraLite®594 Anti-Human NRF2, NFE2L2 (CL594-16396) (red), or 0.4 ug Control Antibody. Cells were fixed and permeabilized with Transcription Factor Staining Buffer Kit (PF00011).