For Research Use Only

CoraLite®594-conjugated Cytochrome c Monoclonal antibody



Catalog Number: CL594-66264

Basic Information

Catalog Number:

 CL594-66264
 BC009578

 Size:
 GeneID (NCBI):

 1000 μ g/ml
 54205

 Source:
 UNIPROT ID:

 Mouse
 Propage

Mouse P99999
Isotype: Full Name: IgG2a cytochrome

IgG2a cytochrome c, somatic
Immunogen Catalog Number: Calculated MW:
AG24349 12 kDa

Purification Method: Protein A purification

CloneNo.: 2D8D11

Excitation/Emission maxima

wavelengths: 588 nm / 604 nm

Applications

Tested Applications:

FC (Intra)

Species Specificity: human, mouse, rat

Background Information

Cytochrome c is a 12-15 kDa electron transporting protein located in the inner mitochondrial membrane. Upon apoptotic stimulation, cytochrome c can be released from mitochondria into cytoplasm, resulting in caspase-3 activation and apoptosis. Measurement of cytochrome c release from the mitochondria is useful for detection of the onset of apoptosis in cells. In addition, cytochrome c can also leave cells and be detectable in extra-cellular medium of apoptotic cells and serum of cancer patients. The level of serum cytochrome c may serve as a prognostic maker during cancer therapy.

GenBank Accession Number:

Storage


Storage:

Store at -20°C. Avoid exposure to light. Stable for one year after shipment. Storage Ruffer

PBS with 50% Glycerol, 0.05% Proclin300, 0.5% BSA, pH 7.3.

Aliquoting is unnecessary for -20°C storage

Selected Validation Data



1X10^6 HepG2 cells were intracellularly stained with 0.4 ug CoraLite®594 Anti-Human Cytochrome c (CL594-66264, Clone:2D8D11) (red), or 0.4 ug Mouse IgG2a Isotype Control (CL594-66360-2, Clone: K11A1B2A2) (blue). Cells were fixed with 4% PFA and permeabilized with Flow Cytometry Perm Buffer (PF00011-C).