For Research Use Only

CoraLite®594-conjugated c-MYC Monoclonal antibody



Catalog Number: CL594-67447

Basic Information

Catalog Number: CL594-67447

Size: 1000 µg/ml Source: Mouse Isotype: lgG1

Immunogen Catalog Number:

AG17519

Calculated MW: 49 kDa

BC000141

4609

P01106

Full Name:

GeneID (NCBI):

UNIPROT ID:

GenBank Accession Number:

v-myc myelocytomatosis viral oncogene homolog (avian)

Observed MW: 55-60 kDa

Purification Method:

Protein G purification

CloneNo.: 3D9C12

Excitation/Emission maxima

wavelengths: 588 nm / 604 nm

Applications

Tested Applications:

FC (Intra)

Species Specificity: Human, mouse, rat

Background Information

MYC contains one basic helix-loop-helix (bHLH) domain. This protein is a multifunctional, nuclear phosphoprotein that plays a role in cell cycle progression, apoptosis and cellular transformation. It functions as a transcription factor $that \ regulates \ transcription \ of \ specific \ target \ genes. \ It \ seems \ to \ activate \ the \ transcription \ of \ growth-related \ genes. \ It \ seems \ to \ activate \ the \ transcription \ of \ growth-related \ genes. \ It \ seems \ to \ activate \ the \ transcription \ of \ growth-related \ genes. \ It \ seems \ to \ activate \ the \ transcription \ of \ growth-related \ genes. \ It \ seems \ to \ activate \ the \ transcription \ of \ growth-related \ genes. \ It \ seems \ to \ activate \ the \ transcription \ of \ growth-related \ genes. \ It \ seems \ to \ activate \ the \ transcription \ of \ growth-related \ genes. \ It \ seems \ to \ activate \ the \ transcription \ of \ growth-related \ genes. \ It \ seems \ to \ activate \ the \ transcription \ of \ growth-related \ genes. \ It \ seems \ to \ activate \ the \ transcription \ of \ growth-related \ genes. \ It \ seems \ the \ transcription \ of \ growth-related \ genes. \ It \ seems \ the \ transcription \ of \ growth-related \ genes. \ the \ transcription \ growth-related \ genes. \ The \ transcription \ the \ transcription \ growth-related \ growth-related \ genes. \ The \ transcription \ the \ tr$ binds DNA in a non-specific manner, yet also specifically recognizes the core sequence 5'-CAC[GA]TG-3'. Mutations, overexpression, rearrangement and translocation of this gene have been associated with a variety of hematopoietic tumors, leukemias and lymphomas, including Burkitt lymphoma.

Storage


Storage:

Store at -20°C. Avoid exposure to light. Stable for one year after shipment.

PBS with 50% Glycerol, 0.05% Proclin300, 0.5% BSA, pH 7.3.

Aliquoting is unnecessary for -20°C storage

Selected Validation Data



1X10^6 HeLa cells were intracellularly stained with 0.4 ug CoraLite® 594 Anti-Human c-MYC (CL594-67447, Clone:3D9C12) (red), or 0.4 ug Control Antibody. Cells were fixed and permeabilized with Transcription Factor Staining Buffer Kit (PF00011).