

For Research Use Only

CoraLite® Plus 647-conjugated EIF4A2 Monoclonal antibody



Catalog Number:CL647-67799

Basic Information

Catalog Number:

CL647-67799

Size:

1 mg/mL

Source:

Mouse

Isotype:

IgG2a

Immunogen Catalog Number:

AG9085

GenBank Accession Number:

BC012547

GeneID (NCBI):

1974

UNIPROT ID:

Q14240

Full Name:

eukaryotic translation initiation
factor 4A, isoform 2

Calculated MW:

408 aa, 46 kDa

Observed MW:

46 kDa

Purification Method:

Protein A purification

CloneNo.:

2D11E11

Excitation/Emission maxima
wavelengths:

654 nm / 674 nm

Applications

Tested Applications:

FC (Intra)

Species Specificity:

Human, Mouse, Rat

Background Information

Eukaryotic initiation factor 4A (EIF4A) has an essential role in the binding of mRNA to the 43S preinitiation complex when protein synthesis begins, as EIF4A unwinds RNA secondary structures in the 5'-UTR of mRNAs which is necessary to allow efficient binding of the small ribosomal subunit, and subsequent scanning for the initiator codon. EIF4A2 is a ATP-dependent RNA helicase, which is a subunit of the eIF4F complex involved in cap recognition and is required for mRNA binding to ribosome.

Storage

Storage:

Store at -20°C. Avoid exposure to light. Stable for one year after shipment.

Storage Buffer:

PBS with 50% Glycerol, 0.05% Proclin300, 0.5% BSA, pH 7.3.

Aliquoting is unnecessary for -20°C storage

For technical support and original validation data for this product please contact:


T: 4006900926

E: Proteintech-CN@ptglab.com


W: ptgcn.com

This product is exclusively available under Proteintech Group brand and is not available to purchase from any other manufacturer.

Selected Validation Data



1×10^6 HeLa cells were intracellularly stained with 0.4 μ g CoraLite® Plus 647 Anti-Human EIF4A2 (CL647-67799, Clone:2D11E11) (red), or 0.4 μ g CoraLite® Plus 647 Mouse IgG2a Isotype Control (C1.18.4) (CL647-65208, Clone: C1.18.4) (blue). Cells were fixed with 4% PFA and permeabilized with Flow Cytometry Perm Buffer (PF00011-C).



1×10^6 HEK-293 cells were intracellularly stained with 0.4 μ g CoraLite® Plus 647 Anti-Human EIF4A2 (CL647-67799, Clone:2D11E11) (red), or 0.4 μ g CoraLite® Plus 647 Mouse IgG2a Isotype Control (C1.18.4) (CL647-65208, Clone: C1.18.4) (blue). Cells were fixed with 4% PFA and permeabilized with Flow Cytometry Perm Buffer (PF00011-C).