

# IHC*easy* YAP1 Ready-To-Use IHC Kit

Catalog Number: **KHC1608**

## General Information

**Sample type:**  
FFPE tissue

**Cited sample type:**

**Reactivity:**  
Human, Mouse, Rat

**Cited Reactivity:**

**Assay type:**  
Immunohistochemistry

**Primary antibody type:**  
Rabbit Polyclonal

**Secondary antibody type:**  
Polymer-HRP-Goat anti-Rabbit

## Kit Component

Component	Size	Concentration
Antigen Retrieval Buffer	100 mL	50×
Washing Buffer	100 mL ×2	20×
Blocking Buffer	5 mL	RTU
Primary Antibody	5 mL	RTU
Secondary Antibody	5 mL	RTU
Chromogen Component A	0.2 mL	RTU
Chromogen Component B	4 mL	RTU
Signal Enhancer	5 mL	RTU
Counter Staining Reagent	5 mL	RTU
Mounting Media	5 mL	RTU
Control Slide	1 slide (Optional)	FFPE
Datasheet	1 Copy	
Manual	1 Copy	

## Storage Instructions

All the reagents are stored at 2-8°C. The kit is stable for 6 months from the date of receipt.


## Background

Yes-associated protein 1 (YAP1) is a transcriptional regulator which can act both as a coactivator and a corepressor and is the critical downstream regulatory target in the Hippo signaling pathway that plays a pivotal role in organ size control and tumor suppression by restricting proliferation and promoting apoptosis. The core of this pathway is composed of a kinase cascade wherein STK3/MST2 and STK4/MST1, in complex with its regulatory protein SAV1, phosphorylates and activates LATS1/2 in complex with its regulatory protein MOB1, which in turn phosphorylates and inactivates YAP1 oncoprotein and WWTR1/TAZ. Plays a key role to control cell proliferation in response to cell contact. Phosphorylation of YAP1 by LATS1/2 inhibits its translocation into the nucleus to regulate cellular genes important for cell proliferation, cell death, and cell migration. The presence of TEAD transcription factors are required for it to stimulate gene expression, cell growth, anchorage-independent growth, and epithelial mesenchymal transition (EMT) induction. Isoform 2 and isoform 3 can activate the C-terminal fragment (CTF) of ERBB4 (isoform 3). Increased expression seen in some liver and prostate cancers. Isoforms lacking the transactivation domain found in striatal neurons of patients with Huntington disease. It is activated by phosphorylation and degraded by ubiquitination.


## Synonyms

65 kDa Yes associated protein, YAP, YAP1, YAP2, YAP65, YKI, Yorkie homolog


## Selected Validation Data




Immunohistochemical analysis of paraffin-embedded human cervical cancer tissue slide using KHC1608 (YAP1 IHC Kit).




Immunohistochemical analysis of paraffin-embedded human stomach cancer tissue slide using KHC1608 (YAP1 IHC Kit).




Immunohistochemical analysis of paraffin-embedded mouse lung tissue slide using KHC1608 (YAP1 IHC Kit).




Immunohistochemical analysis of paraffin-embedded mouse ovary tissue slide using KHC1608 (YAP1 IHC Kit).



Immunohistochemical analysis of paraffin-embedded mouse small intestine tissue slide using KHC1608 (YAP1 IHC Kit).



Immunohistochemical analysis of paraffin-embedded rat lung tissue slide using KHC1608 (YAP1 IHC Kit).



Immunohistochemical analysis of paraffin-embedded rat ovary tissue slide using KHC1608 (YAP1 IHC Kit).