

Human CD16 Magnetic Beads Kit

Catalog Number:KMS006

Description

CD16, which is also known as Fc γ RIII, is a receptor for the Fc γ portion of IgG and is involved in antibody dependent cellular cytotoxicity. 5%-10% human peripheral blood mononuclear cells (PBMC) express CD16. Human CD16 Magnetic Beads Kit is used for isolation or depletion of human CD16 NK cells/monocytes from PBMC, whole blood, or other sample types. Following incubation with biotinylated human CD16 antibody and Streptavidin magnetic beads, the cell sample is placed on a magnet. CD16+ cells remain attached to magnetic beads after separation and can be used for downstream applications, such as in cell expansion. CD16- cells remain in supernatant and can also be used for further applications.

Components

KMS006-10:
 · MS001-10: 100 μ L 10mg/mL streptavidin magnetic beads
 · MS65090-10: 100 μ L 0.1mg/mL Biotin-CD16 (clone: 3G8)
 KMS006-100:
 · MS001-100: 1mL 10mg/mL streptavidin magnetic beads
 · MS65090-100: 1mL 0.1mg/mL Biotin-CD16 (clone: 3G8)

Package

10test/100test

Storage

2-8°C

Storage buffer

PBS, pH7.4, 0.2% BSA and 0.05% Sodium Azide

Reactivity


Human

Recommend usage

10 μ L Biotin-CD16 antibody and 10 μ L streptavidin beads for 1*10⁷ cells.

Results

Representative example of depletion



Following cell separation, cell suspension was stained with FITC-CD45(F10-89-4) and CL650-CD16(3G8) antibodies. All viable cells are gated in the analysis. Left panel: CD16+CD45+ cells before selection. Right panel: CD16+CD45+ cells after depletion. Human CD16 magnetic beads kit is tested using PBMC from three donors.

For technical support and original validation data for this product please contact:

T: 027-87531629 E: Proteintech-CN@ptglab.com W: ptgcn.com

This product is exclusively available under Proteintech Group brand and is not available to purchase from any other manufacturer.