

## cIAP1 Polyclonal antibody

Catalog Number: 10022-1-AP

Featured Product

19 Publications

## Basic Information

## Catalog Number:

10022-1-AP

## Size:

800 µg/ml

## Source:

Rabbit

## Isotype:

IgG

## Immunogen Catalog Number:

AG15203

## GenBank Accession Number:

BC016174

## GeneID (NCBI):

329

## UNIPROT ID:

Q13490

## Full Name:

baculoviral IAP repeat-containing 2

## Calculated MW:

618 aa, 70 kDa

## Observed MW:

70 kDa, 55-60 kDa

## Purification Method:

Antigen affinity purification

## Recommended Dilutions:

WB 1:500-1:2000

IHC 1:50-1:500

IF 1:200-1:800

## Applications

## Tested Applications:

IF/ICC, IHC, WB, ELISA

## Cited Applications:

IF, IHC, WB

## Species Specificity:

human, mouse

## Cited Species:

human, mouse, zebrafish

**Note-IHC: suggested antigen retrieval with TE buffer pH 9.0; (\*) Alternatively, antigen retrieval may be performed with citrate buffer pH 6.0**

## Positive Controls:

**WB**: mouse skeletal muscle tissue, Jurkat cells, mouse brain tissue, mouse liver tissue, mouse testis tissue

**IHC**: human testis tissue, mouse pancreas tissue, mouse testis tissue, human pancreas tissue

**IF**: HepG2 cells,

## Background Information

BIRC2 (also known as cIAP1) is a member of the inhibitor of apoptosis protein (IAP) family. The inhibitor of apoptosis (IAP) proteins are a family of anti-apoptotic regulators found in viruses and metazoans. BIRC2 is a nuclear shuttling protein, whose subcellular localization is mediated by the CRM1-dependent nuclear export pathway (PMID: 15265700). The protein is regulated transcriptionally and can be inhibited by mitochondrial proteins released in the cytoplasm upon apoptotic stimuli (PMID: 15187025). BIRC2 is also believed to be a critical regulator of vascular integrity and endothelial cell survival, thereby providing an additional target pathway for the control of angiogenesis and blood vessel homeostasis during embryogenesis, regeneration and tumorigenesis (PMID: 17934460). This BIRC2 antibody (10022-1-AP) can bind both full length (70kd) and cleaved form (60kd) of the protein.

## Notable Publications

Author	Pubmed ID	Journal	Application
Lei Wang	32882585	Biomed Pharmacother	WB
Michael Hinz	20932475	Mol Cell	WB
Dongsheng Nie	26607717	Biol Reprod	WB

## Storage

## Storage:

Store at -20°C. Stable for one year after shipment.

## Storage Buffer:

PBS with 0.02% sodium azide and 50% glycerol pH 7.3.

Aliquoting is unnecessary for -20°C storage


For technical support and original validation data for this product please contact:

T: 4006900926


E: [Proteintech-CN@ptglab.com](mailto:Proteintech-CN@ptglab.com)W: [ptgcn.com](http://ptgcn.com)

**This product is exclusively available under Proteintech Group brand and is not available to purchase from any other manufacturer.**


## Selected Validation Data




mouse skeletal muscle tissue were subjected to SDS PAGE followed by western blot with 10022-1-AP (cIAP1 antibody) at dilution of 1:1000 incubated at room temperature for 1.5 hours.




Immunohistochemical analysis of paraffin-embedded human testis tissue slide using 10022-1-AP (cIAP1 Antibody) at dilution of 1:200 (under 10x lens).




Immunohistochemical analysis of paraffin-embedded human testis tissue slide using 10022-1-AP (cIAP1 Antibody) at dilution of 1:200 (under 40x lens).




Immunohistochemical analysis of paraffin-embedded mouse pancreas tissue slide using 10022-1-AP (cIAP1 antibody) at dilution of 1:200 (under 40x lens). Heat mediated antigen retrieval with Tris-EDTA buffer (pH 9.0).



Immunohistochemical analysis of paraffin-embedded mouse testis tissue slide using 10022-1-AP (cIAP1 antibody) at dilution of 1:200 (under 40x lens). Heat mediated antigen retrieval with Tris-EDTA buffer (pH 9.0).



Immunofluorescent analysis of (4% PFA) fixed HepG2 cells using 10022-1-AP (cIAP1 antibody), at dilution of 1:200 and CoraLite®488-Conjugated AffiniPure Goat Anti-Rabbit IgG(H+L).



Immunofluorescent analysis of (-20°C Methanol) fixed HepG2 cells using cIAP1 antibody (10022-1-AP) at dilution of 1:400 and CoraLite®488-Conjugated AffiniPure Goat Anti-Rabbit IgG(H+L).