For Research Use Only

Importin Beta Polyclonal antibody

Catalog Number: 10077-1-AP

Featured Product

11 Publications



Basic Information

Catalog Number:

10077-1-AP

Size:

600 µg/ml

Source:

Rabbit

GenBank Accession Number:

BC003572

GeneID (NCBI):

3837

Surce:

UNIPROT ID:

Rabtit

Q14974

Isotype:

Full Name:

G karyopherin (importin) beta 1

Immunogen Catalog Number: Calculated MW: AG0122 97 kDa

Observed MW: 90-97 kDa

Applications

Tested Applications: FC, IF/ICC, IHC, WB, ELISA Cited Applications:

WB, IP, IHC, CoIP Species Specificity: human, mouse, rat Cited Species: human, mouse

Note-IHC: suggested antigen retrieval with TE buffer pH 9.0; (*) Alternatively, antigen retrieval may be performed with citrate buffer pH 6.0 Positive Controls:

WB: A2780 cells, mouse brain tissue, rat brain tissue, HeLa cells, NIH/3T3 cells, C6 cells, HT-1080 cells, PC-12 cells

Purification Method:

WB 1:1000-1:4000 IHC 1:20-1:200

IF 1:50-1:500

Antigen affinity purification

Recommended Dilutions:

IHC: human colon cancer tissue,

IF: HeLa cells,

Background Information

Importin beta 1, also known karyopherin beta or KPNB1, is a nuclear transport receptor that plays a key role in the nuclear import process. Importin beta 1 and importin alpha work in concert to transport cargo into the nucleus, functioning as an essential component of the classical nuclear localization signal (NLS) pathway. Elevated expression of importin beta 1 has been found in cervical tumors. (22125623)

Notable Publications

Author	Pubmed ID	Journal	Application
Zongfu Pan	36192735	Mol Cancer	WB
Hongmin Yao	36300763	Autophagy	WB,IHC
Anna Nakanishi	36366568	Viruses	WB

Storage

Storage


Store at -20°C. Stable for one year after shipment.

Storage Buffer


PBS with 0.02% sodium azide and 50% glycerol pH 7.3.

Aliquoting is unnecessary for -20°C storage


Selected Validation Data




Various lysates were subjected to SDS PAGE followed by western blot with 10077-1-AP (Importin Beta antibody) at dilution of 1:2000 incubated at room temperature for 1.5 hours.



Immunohistochemical analysis of paraffinembedded human colon cancer using 10077-1-AP (Importin beta antibody) at dilution of 1:50 (under 10x lens).



Immunofluorescent analysis of (4% PFA) fixed HeLa cells using Importin Beta antibody (10077-1-AP) at dilution of 1:200 and CoraLite® 488-Conjugated AffiniPure Goat Anti-Rabbit IgG(H+L), CL594-Phalloidin (red).



1X10^6 Jurkat cells were intracellularly stained with 0.4 ug Anti-Human Importin Beta (10077-1-AP) and Coralite®488-Conjugated AffiniPure Goat Anti-Rabbit IgG(H+L) at dilution 1:1000 (red), or 0.4 ug Isotype Control. Cells were fixed with 4% PFA and permeabilized with Flow Cytometry Perm Buffer (PF00011-C).