For Research Use Only

CX3CL1 Polyclonal antibody Catalog Number:10108-2-AP 14 Publications



Basic Information	Catalog Number: 10108-2-AP	GenBank Accession Number: BC001163		Purification Method: Antigen affinity purification	
	Concentration:	GenelD (NCBI):		Recommended Dilutions:	
	1000 ug/ml Source:	6376 UNIPROT ID: P78423		WB 1:500-1:2000 IHC 1:200-1:800	
	Rabbit				
	lsotype: IgG	ype: Full Name: chemokine (C-X3-C motif) ligand 1			
	Immunogen Catalog Number:Calculated MW:AG016142 kDa				
		Observed MW: 90-100 kDa			
Applications	Tested Applications:		Positive Controls:		
	WB, IHC, ELISA Cited Applications:		WB : PC-3 cells, human brain tissue		
	WB, IHC	IHC : human small intestine tissue, human lung tissue			
	Species Specificity: human				
	Cited Species: human, mouse				
	Note-IHC: suggested antigen retrieval with TE buffer pH 9.0; (*) Alternatively, antigen retrieval may be performed with citrate buffer pH 6.0				
Background Information	Chemokines direct the trafficking of white blood cells in immune surveillance, playing a key role in inflammatory and infectious diseases such as AIDS. The human CX3C chemokine carries the chemokine domain on top of an extended mucin-like stalk. CX3CL1 is synthesized as a 50-75 kDa precursor and undergoes glycosylation to yield two mature forms: a 100 kDa cell membrane bound form and an 85 kDa soluble form which is released from the cell surface. The soluble CX3C chemokine has potent chemoattractant activity for T cells and monocytes, and the cell-surface-bound protein, which is induced by primary proinflammatory signals in activated primary endothelial cells promotes strong adhesion of those leukocytes.				
Notable Publications	Author	Pubmed ID J	ournal	Application	
	Hai-Yan Fu		Cells	WB	
	Pingfu Hou		nt J Biol Sci	WB	
	Shuo Miao	32799418 M	Iol Oncol	WB	
	Storage:				

For technical support and original validation data for this product please contact: E: Proteintech-CN@ptglab.com T: 4006900926 W: ptgcn.com

This product is exclusively available under Proteintech Group brand and is not available to purchase from any other manufacturer.


Selected Validation Data





PC-3 cells were subjected to SDS PAGE followed by western blot with 10108-2-AP (CX3CL1 antibody) at dilution of 1:1000 incubated at room temperature for 1.5 hours.

Immunohistochemical analysis of paraffinembedded human small intestine tissue slide using 10108-2-AP (CX3CL1 antibody) at dilution of 1:400 (under 10x lens)..



Immunohistochemical analysis of paraffinembedded human small intestine tissue slide using 10108-2-AP (CX3CL1 antibody) at dilution of 1:400 (under 40x lens)..