For Research Use Only

CORO2A Polyclonal antibody

Catalog Number:10119-1-AP

Featured Product





Basic Information	Catalog Number: 10119-1-AP	GenBank Accession Number BC000010	: Purification Method: Antigen affinity purification
	Size: 400 µg/ml Source: Rabbit	GeneID (NCBI): 7464 UNIPROT ID: Q92828	Recommended Dilutions: WB 1:1000-1:4000 IHC 1:50-1:500
	Isotype: IgG	Full Name: coronin, actin binding protein, 2A Calculated MW: 60 kDa	
	Immunogen Catalog Number: AG0164		
		Observed MW: 48 kDa	
Applications	human mouse		
			IHC : human skin cancer tissue, mouse small intestine tissue
	Note-IHC: suggested antigen retrieval with TE buffer pH 9.0; (*) Alternatively, antigen retrieval may be performed with citrate buffer pH 6.0		
Background Information	CORO2A (Coronin 2A), also named as CLIPINB, is a member of the 'short' coronin subfamily containing a single WD40-repeat domain. It has been shown that the expression of CORO2A is up-regulated in human colon cancer. CORO2A can interact with MAPK14 and PRMT5. The calculated MW of CORO2A is 60 kDa, while 10119-1-AP antibody detects 48 kDa protein which is similar to the paper published. (PMID: 26373535)		
Storage	Storage: Store at -20°C. Stable for one year Storage Buffer: PBS with 0.02% sodium azide and Aliquoting is unnecessary for -20°	50% glycerol pH 7.3.	

For technical support and original validation data for this product please contact:T: 4006900926E: Proteintech-CN@ptglab.comW: ptgcn.com

This product is exclusively available under Proteintech Group brand and is not available to purchase from any other manufacturer.

Selected Validation Data





Various lysates were subjected to SDS PAGE followed by western blot with 10119-1-AP (CORO2A antibody) at dilution of 1:2000 incubated at room temperature for 1.5 hours. Immunohistochemical analysis of paraffinembedded human skin cancer tissue slide using 10119-1-AP (CORO2A antibody) at dilution of 1:200 (under 10x lens). Heat mediated antigen retrieval with Tris-EDTA buffer (pH 9.0). WB result of CORO2A antibody (10119-1-AP; 1:4000; incubated at room temperature for 1.5 hours) with sh-Control and sh-CORO2A transfected Jurkat cells.