

For Research Use Only

FUSIP1 Polyclonal antibody

Catalog Number: 10131-1-AP **3 Publications**



Basic Information

Catalog Number: 10131-1-AP	GenBank Accession Number: BC001107	Purification Method: Antigen affinity purification
Size: 500 µg/ml	GeneID (NCBI): 10772	Recommended Dilutions: WB 1:500-1:2000 IHC 1:250-1:1000
Source: Rabbit	UNIPROT ID: O75494	
Isotype: IgG	Full Name: FUS interacting protein (serine/arginine-rich) 1	
Immunogen Catalog Number: AG0176	Calculated MW: 22 kDa	
	Observed MW: 31 kDa	

Applications

Tested Applications: IHC, WB, ELISA	Positive Controls:
Cited Applications: WB, IHC	WB: HepG2 cells, HeLa cells
Species Specificity: human, mouse, rat	IHC: human urothelial carcinoma tissue,
Cited Species: human	

Note-IHC: suggested antigen retrieval with TE buffer pH 9.0; (*) Alternatively, antigen retrieval may be performed with citrate buffer pH 6.0

Background Information

FUSIP1, also named as SRp40, SRp38, FUSIP2, SFRS13A, or TASR, is a 262 amino acid protein, which belongs to the splicing factor SR family. FUSIP1 as a Splicing factor that in its dephosphorylated form acts as a general repressor of pre-mRNA splicing. FUSIP1 is required for splicing repression in M-phase cells and after heat shock and also acts as a splicing factor that specifically promotes exon skipping during alternative splicing. FUSIP1 Interacting with YTHDC1, a RNA-binding protein that recognizes and binds N6-methyladenosine (m6A)-containing RNAs, prevents SRSF10 from binding to its mRNA-binding sites close to m6A-containing regions, leading to inhibit exon skipping during alternative splicing.

Notable Publications

Author	Pubmed ID	Journal	Application
Xiaobai Liu	32600379	J Exp Clin Cancer Res	WB
Yuhan Chen	29942010	Nat Commun	WB
Yi-Ting Chen	38196148	Proteomics Clin Appl	WB,IHC

Storage

Storage:
Store at -20°C. Stable for one year after shipment.
Storage Buffer:
PBS with 0.02% sodium azide and 50% glycerol pH 7.3.
Aliquoting is unnecessary for -20°C storage

For technical support and original validation data for this product please contact:


T: 4006900926

E: Proteintech-CN@ptglab.com


W: ptgcn.com

This product is exclusively available under Proteintech Group brand and is not available to purchase from any other manufacturer.


Selected Validation Data



HepG2 cells were subjected to SDS PAGE followed by western blot with 10131-1-AP (FUSIP1 Antibody) at dilution of 1:1000 incubated at room temperature for 1.5 hours.



Immunohistochemical analysis of paraffin-embedded human urothelial carcinoma tissue slide using 10131-1-AP (FUSIP1 antibody) at dilution of 1:500 (under 10x lens). Heat mediated antigen retrieval with Tris-EDTA buffer (pH 9.0).



Immunohistochemical analysis of paraffin-embedded human urothelial carcinoma tissue slide using 10131-1-AP (FUSIP1 antibody) at dilution of 1:500 (under 40x lens). Heat mediated antigen retrieval with Tris-EDTA buffer (pH 9.0).