

For Research Use Only

# ATG5 Polyclonal antibody

Catalog Number: 10181-2-AP

Featured Product

338 Publications



## Basic Information

**Catalog Number:**

10181-2-AP

**Concentration:**

950 ug/ml

**Source:**

Rabbit

**Isotype:**

IgG

**Immunogen Catalog Number:**

AG0214

**GenBank Accession Number:**

BC002699

**GeneID (NCBI):**

9474

**UNIPROT ID:**

Q9H1Y0

**Full Name:**

ATG5 autophagy related 5 homolog  
(S. cerevisiae)

**Calculated MW:**

32 kDa

**Observed MW:**

32 kDa, 40-45 kDa, 50-55 kDa

**Purification Method:**

Antigen affinity purification

**Recommended Dilutions:**

WB 1:1000-1:5000

IP 0.5-4.0 ug for 1.0-3.0 mg of total protein lysate

## Applications

**Tested Applications:**

WB, IP, ELISA

**Cited Applications:**

WB, IHC, IF, IP, CoIP

**Species Specificity:**

human, mouse, rat

**Cited Species:**

human, mouse, rat, pig, monkey, chicken, bovine, hamster, goat, grouper

**Positive Controls:**

WB: A549 cells, HeLa cells, mouse kidney tissue, HepG2 cells

IP: HeLa cells,

## Background Information

ATG5, also named as APG5L and ASP, belongs to the ATG5 family. It is required for autophagy. It plays an important role in the apoptotic process, possibly within the modified cytoskeleton. Its expression is a relatively late event in the apoptotic process, occurring downstream of caspase activity. Autophagy is a catabolic process for the autophagosomic-lysosomal degradation of bulk cytoplasmic contents. Formation of the autophagosome involves a ubiquitin-like conjugation system in which Atg12 is covalently bound to Atg5 and targeted to autophagosome vesicles. It mediates autophagosome-independent host protection. This antibody is raised against 28-275 amino acids of human ATG5. It can recognize the ATG5-ATG12 complex (55 kDa) which can be truncated and generate a 40-45 kDa band. 10181-2-AP also recognizes the free ATG5 (32 kDa).

## Notable Publications

Author	Pubmed ID	Journal	Application
Zeen Zhu	36248959	Front Oncol	WB
Samana Batool	30274346	Int J Mol Sci	WB
Jiawei Hao	36126167	Autophagy	WB

## Storage

**Storage:**

Store at -20°C. Stable for one year after shipment.

**Storage Buffer:**

PBS with 0.02% sodium azide and 50% glycerol pH 7.3.

Aliquoting is unnecessary for -20°C storage

For technical support and original validation data for this product please contact:


T: 4006900926

E: Proteintech-CN@ptglab.com


W: ptgcn.com

This product is exclusively available under Proteintech Group brand and is not available to purchase from any other manufacturer.


## Selected Validation Data




HeLa cells were subjected to SDS PAGE followed by western blot with 10181-2-AP (ATG5 antibody) at dilution of 1:500 incubated at room temperature for 1.5 hours.



Various lysates were subjected to SDS PAGE followed by western blot with 10181-2-AP (ATG5 antibody) at dilution of 1:2500 incubated at room temperature for 1.5 hours.



WB result of ATG5 antibody (10181-2-AP; 1:2000; room temperature for 1.5 hours) with wild-type and ATG5 knockout HeLa cells.



IP result of anti-ATG5 (IP:10181-2-AP, 4ug; Detection:10181-2-AP 1:3000) with HeLa cells lysate 1760 ug.