For Research Use Only

TSC22D1 Polyclonal antibody

Catalog Number:10214-1-lg Featured Product 9 Publications





Basic Information	Catalog Number: 10214-1-lg	GenBank Accession Number: BC000456	Purification Method: Protein A purification
	Concentration:	GeneID (NCBI):	Recommended Dilutions:
	1000 ug/ml	8848	WB 1:500-1:1000
	Source:	UNIPROT ID:	IP 0.5-4.0 ug for 1.0-3.0 mg of total protein lysate
	Rabbit	Q15714 Full Name:	IHC 1:20-1:200
	lsotype: IgG	TSC22 domain family, member 1	IF/ICC 1:200-1:800
	Immunogen Catalog Number: AG0287	Calculated MW: 16 kDa	
		Observed MW: 16 kDa	
Applications	Tested Applications: Positive Controls:		
	WB, IHC, IF/ICC, IP, ELISA WB : human Cited Applications:		brain tissue, A549 cells, rat brain tissue
	WB, IHC, IF, IP	IP : rat brain	tissue,
	Species Specificity:		brain tissue, human testis tissue
	human, rat	IF/ICC : A54	9 cells,
	Cited Species: human, mouse		
	Note-IHC: suggested antigen retrieval with TE buffer pH 9.0; (*) Alternatively, antigen retrieval may be performed with citrate buffer pH 6.0		
Background Information	TGFB114, MGC 17597) is a membe activity and belongs to the large f and cell death. Leucine zipper stru is also an important downstream	ning growth factor beta 1 induced transc r of a family of leucine zipper containing amily of early response genes. TSC-22 r icture of TSC22 inhibits the anchorage ir component of peroxisome proliferator-a lial cell differentiation. This antibody is an TSC22D1.	g transcription factors with repressor egulates cell growth and differentiatio idependent growth of cancer cells. TSC- ctivated receptor (PPAR) and TGF-beta
	TGFB114, MGC 17597) is a membe activity and belongs to the large f and cell death. Leucine zipper stru is also an important downstream signaling during intestinal epithe	r of a family of leucine zipper containing amily of early response genes. TSC-22 r icture of TSC22 inhibits the anchorage ir component of peroxisome proliferator-a lial cell differentiation. This antibody is	g transcription factors with repressor egulates cell growth and differentiatio idependent growth of cancer cells. TSC- ctivated receptor (PPAR) and TGF-beta
	TGFB114, MGC 17597) is a membe activity and belongs to the large f and cell death. Leucine zipper stru is also an important downstream signaling during intestinal epithe against an internal region of hum	r of a family of leucine zipper containing amily of early response genes. TSC-22 r icture of TSC22 inhibits the anchorage ir component of peroxisome proliferator-a lial cell differentiation. This antibody is an TSC22D1.	g transcription factors with repressor egulates cell growth and differentiation dependent growth of cancer cells. TSC- ctivated receptor (PPAR) and TGF-beta a rabbit polyclonal antibody raised
	TGFB114, MGC 17597) is a membe activity and belongs to the large f and cell death. Leucine zipper stru is also an important downstream signaling during intestinal epithe against an internal region of hum Author	r of a family of leucine zipper containing amily of early response genes. TSC-22 r icture of TSC22 inhibits the anchorage ir component of peroxisome proliferator-a lial cell differentiation. This antibody is an TSC22D1. Pubmed ID Journal	g transcription factors with repressor egulates cell growth and differentiation idependent growth of cancer cells. TSC- ctivated receptor (PPAR) and TGF-beta a rabbit polyclonal antibody raised Application
	TGFB114, MGC 17597) is a membe activity and belongs to the large f and cell death. Leucine zipper stru is also an important downstream signaling during intestinal epithe against an internal region of hum Author Ryouta Kamimura	r of a family of leucine zipper containing family of early response genes. TSC-22 r incure of TSC22 inhibits the anchorage in component of peroxisome proliferator-a lial cell differentiation. This antibody is an TSC22D1. Pubmed ID Journal 34681573 Int J Mol Sci	g transcription factors with repressor egulates cell growth and differentiation idependent growth of cancer cells. TSC- ctivated receptor (PPAR) and TGF-beta a rabbit polyclonal antibody raised Application WB,IF WB
Background Information	TGFB114, MGC 17597) is a membe activity and belongs to the large f and cell death. Leucine zipper stru is also an important downstream signaling during intestinal epithe against an internal region of hum Author Ryouta Kamimura Zhi Zheng	r of a family of leucine zipper containing amily of early response genes. TSC-22 r icture of TSC22 inhibits the anchorage ir component of peroxisome proliferator-a lial cell differentiation. This antibody is an TSC22D1. Pubmed ID Journal 34681573 Int J Mol Sci 35513372 Cell Death Discov	g transcription factors with repressor egulates cell growth and differentiatio idependent growth of cancer cells. TSC- ctivated receptor (PPAR) and TGF-beta a rabbit polyclonal antibody raised Application WB,IF WB


For technical support and original validation data for this product please contact: E: Proteintech-CN@ptglab.com T: 4006900926 W: ptgcn.com

This product is exclusively available under Proteintech Group brand and is not available to purchase from any other manufacturer.


Selected Validation Data









IP result of anti-TSC 22D1 (IP:10214-1-Ig, 4ug; Detection:10214-1-Ig 1:500) with rat brain tissue lysate 4000ug.



Immunohistochemical analysis of paraffinembedded human brain tissue slide using 10214-1-Ig (TSC22D1 Antibody) at dilution of 1:50 (under 40x lens).



Immunohistochemical analysis of paraffinembedded human testis tissue slide using 10214-1-Ig (TSC22D1 Antibody) at dilution of 1:50 (under 40x lens).



Immunofluorescent analysis of (4% PFA) fixed A549 cells using TSC22D1 antibody (10214-1-Ig) at dilution of 1:400 and CoraLite®488-Conjugated Goat Anti-Rabbit IgG(H+L) (SA00013-2), CL594phalloidin (red).