For Research Use Only

CDC37 Polyclonal antibody, PBS Only

Catalog Number:10218-1-PBS Featured Product



Basic Information

Catalog Number:

GenBank Accession Number:

Purification Method: Antigen affinity purification

10218-1-PBS

GeneID (NCBI): 11140

BC000083

1 mg/ml Source:

Size:

UNIPROT ID: Q16543 Full Name:

Rabbit Isotype: IgG

cell division cycle 37 homolog (S.

Immunogen Catalog Number:

cerevisiae)
Calculated MW:

AG0267

44 kDa
Observed MW:
45-50 kDa

Applications

Tested Applications:

WB, IHC, IF/ICC, FC (Intra), IP, Indirect ELISA

Species Specificity: human, mouse, monkey

Background Information

Cdc (cell division control) family of proteins are regulatory proteins involved in the processes of cell cycle and proliferation. CDC37, migrated as a protein of 50 kDa in SDS PAGE, is a co-chaperone protein that recruits several immature client kinases to Hsp90 for proper folding. It functions in association with Hsp90, although little is known of its mechanism of action or where it fits into a folding pathway involving other Hsp90 cochaperones. CDC37 upregulation is a common early event in some localized human cancers. But, the effect of CDC37 dysregulation in human cells is unclear.

Storage


Storage:

Store at -80°C.


The product is shipped with ice packs. Upon receipt, store it immediately at -80°C $\,$

Storage Buffer: PBS Only


Selected Validation Data




HEK-293 cells were subjected to SDS PAGE followed by western blot with 10218-1-AP (CDC37 antibody) at dilution of 1:1000 incubated at room temperature for 1.5 hours. This data was developed using the same antibody clone with 10218-1-PBS in a different storage buffer formulation.




Immunohistochemical analysis of paraffinembedded human colon cancer tissue slide using 10218-1-AP (CDC37 Antibody) at dlution of 1:50 (under 10x lens). This data was developed using the same antibody clone with 10218-1-PBS in a different storage buffer formulation.




Immunohistochemical analysis of paraffinembedded human colon cancer tissue slide using 10218-1-AP (CDC 37 Antibody) at dilution of 1:50 (under 40x lens). This data was developed using the same antibody clone with 10218-1-PBS in a different storage buffer formulation.



IP result of anti-CDC37 (IP:10218-1-AP, 4ug; Detection:10218-1-AP 1:1000) with HEK-293 cells lysate 3200ug. This data was developed using the same antibody clone with 10218-1-PBS in a different storage buffer formulation.



1x10^6 MCF-7 cells were intracellularly stained with 0.8 ug CDC37 Polyclonal antibody (10218-1-AP) and CoraLite® 488-Conjugated AffiniPure Goat Anti-Rabbit 1gG(H+L) (SA00013-2)(red), or 0.8 ug Rabbit 1gG control Rabbit PolyAb (30000-0-AP) (blue). Cells were fixed with 4% PFA and permeabilized with Flow Cytometry Perm Buffer (PF00011-C). This data was developed using the same antibody clone with 10218-1-PBS in a different storage buffer formulation.



Immunofluorescent analysis of (-20°C Ethanol) fixed HeLa cells using CDC37 antibody (10218-1-AP) at dilution of 1:400 and Coralite®594-Conjugated Goat Anti-Rabbit IgG(H+L) (SA00013-4). This data was developed using the same antibody clone with 10218-1-PBS in a different storage buffer formulation