For Research Use Only

COPS6 Polyclonal antibody

Catalog Number: 10256-1-AP

1 Publications



Basic Information

Catalog Number: GenBank Accession Number: 10256-1-AP BC002520 GeneID (NCBI): Size: $600 \mu g/ml$ 10980 **UNIPROT ID:** Source: Rabbit Q7L5N1 Full Name: Isotype:

COP9 constitutive photomorphogenic homolog subunit 6 (Arabidopsis)

Immunogen Catalog Number:

AG0331

Calculated MW: 327 aa, 36 kDa Observed MW: 34 kDa

Applications

Tested Applications: IHC, WB, ELISA **Cited Applications:**

Species Specificity:

human, mouse, rat, monkey, pig

Cited Species:

Note-IHC: suggested antigen retrieval with TE buffer pH 9.0; (*) Alternatively, antigen retrieval may be performed with citrate

buffer pH 6.0

Antigen affinity purification Recommended Dilutions:

WB 1:500-1:3000 IHC 1:50-1:500

Purification Method:

Positive Controls:

WB: mouse colon tissue, 3T3-L1 cells, pig lymph node

IHC: human colon cancer tissue,

Background Information

The COPS6 protein, also named as mov34, is a 34kd subunit of COP9 signal osome, a highly conserved eight-subunit $protein\ complex\ that\ functions\ as\ an\ important\ regulator\ in\ multiple\ signaling\ pathways.\ This\ protein\ belongs\ to\ protein\ protein$ translation initiation factor 3 (eIF3) superfamily. It is involved in the regulation of cell cycle and acts as the regulatory subunit of the 26S proteasome.

Notable Publications

| Author | Pubmed ID | Journal | Application |
|---------|-----------|----------------|-------------|
| Zun Mao | 31223289 | Int J Biol Sci | WB |

Storage


Storage:

Store at -20°C. Stable for one year after shipment.


PBS with 0.02% sodium azide and 50% glycerol pH 7.3.

Aliquoting is unnecessary for -20°C storage

Selected Validation Data



Immunohistochemical analysis of paraffinembedded human colon cancer using 10256-1-AP (COPS6 antibody) at dilution of 1:50 (under 10x lens).



Various lysates were subjected to SDS PAGE followed by western blot with 10256-1-AP (COPS6 antibody) at dilution of 1:1500 incubated at room temperature for 1.5 hours.



3T3-L1 cells were subjected to SDS PAGE followed by western blot with 10256-1-AP (COPS6 antibody) at dilution of 1:600 incubated at room temperature for 1.5 hours.