For Research Use Only

H3F3B Polyclonal antibody

Catalog Number: 10265-1-AP 2 Publications



Basic Information

Catalog Number: 10265-1-AP

GenBank Accession Number: BC001124

Purification Method: Antigen affinity purification

Size:

GeneID (NCBI):

300 ug/ml

3021

Recommended Dilutions: WB 1:200-1:1000

Source:

UNIPROT ID: P84243

Full Name:

Rabbit Isotype:

AG0326

H3 histone, family 3B (H3.3B)

Immunogen Catalog Number:

Calculated MW:

15 kDa

Observed MW:

17 kDa

Applications

Tested Applications:

WB, ELISA

WB: HEK-293 cells, HeLa cells

Positive Controls:

Cited Applications: WB

Species Specificity:

human

Cited Species:

human, mouse

Background Information

Histones are the basic nuclear proteins responsible for the nucleosome structure of the chromosomal fiber. Five classes of histone genes have been reported, some of which are expressed only during S phase, while others are replication independent. The latter are referred to as replacement histones and are expressed in quiescent or terminally differentiated cells. H3F3B is one replacement histone [PMID: 8586426]. It constitutes the predominant form of histone H3 in non-dividing cells and is incorporated into chromatin independently of DNA synthesis. Deposited at sites of nucleosomal displacement throughout transcribed genes, suggesting that it represents an epigenetic imprint of transcriptionally active chromatin. Nucleosomes wrap and compact DNA into chromatin, limiting DNA accessibility to the cellular machineries which require DNA as a template. Thereby it has a central role in transcription regulation, DNA repair, DNA replication and chromosomal stability [PMID: 14718166, 15776021].

Notable Publications

Author	Pubmed ID	Journal	Application
Jiajia Wang	38898473	Cell Commun Signal	WB
Bo Dong	38997894	Immunobiology	WB

Storage


Store at -20°C. Stable for one year after shipment.

Storage Buffer:

PBS with 0.02% sodium azide and 50% glycerol pH 7.3.

Aliquoting is unnecessary for -20°C storage

Selected Validation Data



HEK-293 cells were subjected to SDS PAGE followed by western blot with 10265-1-AP (H3F3B antibody) at dilution of 1:300 incubated at room temperature for 1.5 hours.