

For Research Use Only

# CYPC Polyclonal antibody

Catalog Number: 10287-2-AP

Featured Product

7 Publications



## Basic Information

Catalog Number:

10287-2-AP

Size:

600 µg/ml

Source:

Rabbit

Isotype:

IgG

Immunogen Catalog Number:

AG0209

GenBank Accession Number:

BC002678

GeneID (NCBI):

5480

UNIPROT ID:

P45877

Full Name:

peptidylprolyl isomerase C  
(cyclophilin C)

Calculated MW:

23 kDa

Observed MW:

23 kDa

Purification Method:

Antigen affinity purification

Recommended Dilutions:

WB 1:500-1:3000

IHC 1:20-1:200

IF/ICC 1:200-1:800

## Applications

Tested Applications:

WB, IHC, IF/ICC, ELISA

Cited Applications:

WB, IHC

Species Specificity:

human, mouse, rat

Cited Species:

human, mouse

**Note-IHC: suggested antigen retrieval with  
TE buffer pH 9.0; (\*) Alternatively, antigen  
retrieval may be performed with citrate  
buffer pH 6.0**

Positive Controls:

WB : HeLa cells, HuH-7 cells, HEK-293 cells, mouse  
heart tissue

IHC : human heart tissue, mouse kidney tissue

IF/ICC : HEK-293 cells,

## Background Information

Cyclophilin C also named as peptidyl-prolyl cis-trans isomerase C (PPIC), is expressed in several bone marrow (BM) hematopoietic progenitors and in T - cell precursors (PMID: 33864384). CYPC/PPIC catalyze the cis-trans isomerization of proline imidic peptide bonds in oligopeptides and accelerate the folding of proteins. Similar to other PPIases, this protein can bind immunosuppressant cyclosporin A. In addition, CYPC/PPIC is involved in many biological processes, including mitochondrial metabolism, REDOX and inflammation, as well as related diseases and conditions, such as ischemia-reperfusion injury, AIDS and cancer.

## Notable Publications

Author	Pubmed ID	Journal	Application
Rebeca Alvaríño	36277217	Front Physiol	WB
Paloma Gómez-Fernández	31108847	Int J Mol Sci	WB
Udayan Chatterji	19380579	J Biol Chem	WB

## Storage

Storage:

Store at -20°C. Stable for one year after shipment.

Storage Buffer:

PBS with 0.02% sodium azide and 50% glycerol pH 7.3.

Aliquoting is unnecessary for -20°C storage

For technical support and original validation data for this product please contact:


T: 4006900926

E: [Proteintech-CN@ptglab.com](mailto:Proteintech-CN@ptglab.com)


W: [ptgcn.com](http://ptgcn.com)

**This product is exclusively available under Proteintech Group brand and is not available to purchase from any other manufacturer.**


## Selected Validation Data




Various lysates were subjected to SDS PAGE followed by western blot with 10287-2-AP (CYPC antibody) at dilution of 1:1500 incubated at room temperature for 1.5 hours.



Immunofluorescent analysis of (-20°C Ethanol) fixed HEK-293 cells using CYPC antibody (10287-2-AP) at dilution of 1:400 and CoraLite® 488-Conjugated Goat Anti-Rabbit IgG(H+L), CL594-Phalloidin (red).



Immunohistochemical analysis of paraffin-embedded human heart tissue slide using 10287-2-AP (CYPC Antibody) at dilution of 1:50 (under 10x lens).



Immunohistochemical analysis of paraffin-embedded human heart tissue slide using 10287-2-AP (CYPC Antibody) at dilution of 1:50 (under 40x lens).