## For Research Use Only

# RARA Polyclonal antibody

Catalog Number: 10331-1-AP 10 Publications



**Basic Information** 

 Catalog Number:
 GenBank Accession Number:

 10331-1-AP
 BC008727

 Size:
 GeneID (NCBI):

 300 ug/ml
 5914

Source: UNIPROT ID: Rabbit P10276
Isotype: Full Name:

IgG retinoic acid receptor, alpha

Immunogen Catalog Number: Calculated MW: AG0420 51 kDa

Observed MW: 50-60 kDa Purification Method: Antigen affinity purification Recommended Dilutions:

WB 1:500-1:1000 IP 0.5-4.0 ug for 1.0-3.0 mg of total

protein lysate IHC 1:50-1:500

**Applications** 

Tested Applications: WB, IHC, IP, ELISA Cited Applications:

WB, IHC, IF, CoIP Species Specificity: human, mouse, rat Cited Species: human, mouse

Note-IHC: suggested antigen retrieval with TE buffer pH 9.0; (\*) Alternatively, antigen retrieval may be performed with citrate buffer pH 6.0 Positive Controls:

WB: K-562 cells, MCF-7 cells

IP: HeLa cells,

IHC : human breast cancer tissue, human testis tissue,

human colon cancer tissue

# **Background Information**

Retinoic acid receptors (RARs) function as nuclear receptors that control gene expression in response to binding of the ligand retinoic acid (RA). Retinoic acid receptor alpha (RAR $^{\,\alpha}$ ), also known as NR1B1 (nuclear receptor subfamily 1, group B, member 1), plays an essential role in the regulation of retinoic acid-induced germ cell development during spermatogenesis. Formation of a complex with histone deacetylases might lead to inhibition of RARE DNA element binding and to transcriptional repression (PubMed:28167758). Transcriptional activation and RARE DNA element binding might be supported by the transcription factor KLF2 (PubMed:28167758).

#### **Notable Publications**

Author	Pubmed ID	Journal	Application
Shyi-Jang Shin	33065162	Metabolism	WB,IHC
Zhen Yu	35278669	Cell Signal	WB
Lu Wang	36482192	Br J Cancer	WB

Storage

Storage:


Store at -20°C. Stable for one year after shipment.

Storage Buffer:


PBS with 0.02% sodium azide and 50% glycerol pH 7.3.

Aliquoting is unnecessary for -20°C storage


## **Selected Validation Data**



K-562 cells were subjected to SDS PAGE followed by western blot with 10331-1-AP (RARA antibody) at dilution of 1:600 incubated at room temperature for 1.5 hours.



Immunohistochemical analysis of paraffinembedded human breast cancer tissue slide using 10331-1-AP (RARA antibody) at dilution of 1:200 (under 40x lens). Heat mediated antigen retrieval with Tris-EDTA buffer (pH 9.0).



IP result of anti-RARA (IP:10331-1-AP, 4ug; Detection:10331-1-AP 1:2000) with HeLa cells lysate 1760 ug.