For Research Use Only

RNH1 Polyclonal antibody Catalog Number: 10345-1-AP Featured Product

Featured Product

12 Publications




Basic Information	Catalog Number: 10345-1-AP	GenBank Accession Number: BC000677		Purification Method: Antigen affinity purification	
	Size:	GenelD (NC	BI):	Recommended Dilutions:	
	700 µg/ml	6050		WB 1:2000-1:10000 IP 0.5-4.0 ug for 1.0-3.0 mg of total	
	Source: Rabbit	UNIPROT ID P13489	t.	protein lysate	
	Isotype:	Full Name:		IHC 1:50-1:500	
	IgG		e/angiogenin inhibitor 1	IF/ICC 1:200-1:800	
	Immunogen Catalog Number: AG0350	Calculated I 50 kDa	Calculated MW: 50 kDa Observed MW: 45-50 kDa		
Applications	Tested Applications:			rols:	
	WB, IHC, IF/ICC, FC (Intra), IP, ELIS	5A		WB : HEK-293T cells, HeLa cells, human brain tissue, human heart tissue, human liver tissue, LO2 cells, HepG2 cells, Jurkat cells, human placenta tissue IP : mouse liver tissue, HEK-293T cells, rat liver tissue IHC : mouse liver tissue, IF/ICC : MCF-7 cells,	
	Cited Applications: WB, IHC, IF				
	Species Specificity:		•		
	human, mouse, rat				
	Cited Species: human, mouse		IF/ICC : MCF-		
	Note-IHC: suggested antigen retrieval with TE buffer pH 9.0; (*) Alternatively, antigen retrieval may be performed with citrate buffer pH 6.0				
Background Information	Ribonuclease/ angiogenin inhibit tightly to ribonucleases in cells an human RNH gene has been region	nd may be essen	tial in the control of mRN	8200, MGC 54054) is a protein that bind A degradation and gene expression. Th s by in situ hybridization.	
Notable Publications	Author	Pubmed ID	Journal	Application	
	Yasutoshi Akiyama	36361884	Int J Mol Sci	WB	
			Front Mol Biosci	WB	
	Yasutoshi Akiyama	35300117	i tonti lot biosei		
	Yasutoshi Akiyama Shozo Honda	26124144	Proc Natl Acad Sci U	S A WB	


For technical support and original validation data for this product please contact: E: Proteintech-CN@ptglab.com T: 4006900926 W: ptgcn.com

This product is exclusively available under Proteintech Group brand and is not available to purchase from any other manufacturer.


Selected Validation Data




Various lysates were subjected to SDS PAGE followed by western blot with 10345-1-AP (RNH1 antibody) at dilution of 1:6000 incubated at room temperature for 1.5 hours.




WB result of RNH1 antibody (10345-1-AP, 1:2500) with si-control and si-RNH1 transfected HeLa cells.




Immunohistochemical analysis of paraffinembedded mouse liver tissue slide using 10345-1-AP (RNH1 antibody) at dilution of 1:200 (under 10x lens). Heat mediated antigen retrieval with Tris-EDTA buffer (pH 9.0).




Immunohistochemical analysis of paraffinembedded mouse liver tissue slide using 10345-1-AP (RNH1 antibody) at dilution of 1:200 (under 40x lens). Heat mediated antigen retrieval with Tris-EDTA buffer (pH 9.0).




Immunofluorescent analysis of (-20°C Methanol) fixed MCF-7 cells using RNH1 antibody (10345-1-AP) at dilution of 1:400 and CoraLite®488-Conjugated Goat Anti-Rabbit IgG(H+L).




IP result of anti-RNH1 (IP:10345-1-AP, 4ug; Detection:10345-1-AP 1:500) with mouse liver tissue lysate 6400ug.




IP result of anti-RNH1 (IP:10345-1-AP, 4ug; Detection:10345-1-AP 1:8000) with HEK-293T cells lysate 2040 ug.



IP result of anti-RNH1 (IP:10345-1-AP, 4ug; Detection:10345-1-AP 1:8000) with mouse liver tissue lysate 1840 ug.



IP result of anti-RNH1 (IP:10345-1-AP, 4ug; Detection:10345-1-AP 1:8000) with rat liver tissue lysate 5520 ug.



1X10^6 MCF-7 cells were intracellularly stained with 0.4 ug Anti-Human RNH1 (10345-1-AP) and CoraLite@488-Conjugated Goat Anti-Rabbit IgG(H+L) at dilution 1:1000 (red), or 0.4 ug Control Antibody. Cells were fixed with 4% PFA and permeabilized with Flow Cytometry Perm Buffer (PF00011-C).