#### For Research Use Only

# GRWD1 Polyclonal antibody

Catalog Number: 10354-1-AP

1 Publications



**Purification Method:** 

WB 1:500-1:1000

protein lysate

IHC 1:20-1:200

Antigen affinity purification

IP 0.5-4.0 ug for 1.0-3.0 mg of total

Recommended Dilutions:

**Basic Information** 

Catalog Number: GenBank Accession Number: 10354-1-AP BC002440
Size: GeneID (NCBI): 83743
Source: UNIPROT ID: Rabbit Q9BQ67
Isotype: Full Name:

glutamate-rich WD repeat containing

Immunogen Catalog Number:

AG0367 Calculated MW:

49 kDa Observed MW: 49 kDa

**Applications** 

Tested Applications: IHC, IP, WB, ELISA Cited Applications: WB, IHC

Species Specificity: human Cited Species:

Note-IHC: suggested antigen retrieval with TE buffer pH 9.0; (\*) Alternatively, antigen retrieval may be performed with citrate

buffer pH 6.0

Positive Controls:

WB: HEK-293 cells, PC-3 cells, SKOV-3 cells

IP: PC-3 cells,

IHC: human prostate cancer tissue,

## **Background Information**

WD-repeat proteins are a class of functionally divergent molecules that cooperate with other proteins to regulate cellular processes. WD-repeat proteins have been implicated in a variety of cellular functions that are mostly regulatory in nature, including cytoskeleton organization, vesicular fusion, pre-mRNA splicing, and cell cycle regulation. WD-repeat-containing protein designated glutamate-rich WD repeat (GRWD1) is a component of the 50S and 80S preribosomal complexes and plays a role in ribosome biogenesis and during myeloid differentiation its levels are regulated. [PMID:15885502]

#### **Notable Publications**

Author	Pubmed ID	Journal	Application
Qiongzi Wang	31891681	Exp Cell Res	WB,IHC

Storage

Storage:


Store at -20°C. Stable for one year after shipment.

Storage Buffer


PBS with 0.02% sodium azide and 50% glycerol pH 7.3.

Aliquoting is unnecessary for -20°C storage


### Selected Validation Data



HEK-293 cells were subjected to SDS PAGE followed by western blot with 10354-1-AP (GRWD1 Antibody) at dilution of 1:600 incubated at room temperature for 1.5 hours.



Immunohistochemical analysis of paraffinembedded human prostate cancer using 10354-1-AP (GRWD1 antibody) at dilution of 1:100 (under 25x Jans)



IP result of anti-GRWD1 (IP:10354-1-AP, 4ug; Detection:10354-1-AP 1:1000) with PC-3 cells lysate 4000ug.