For Research Use Only

RAP1GDS1 Polyclonal antibody

Catalog Number: 10377-1-AP 4 Publications



Basic Information

Catalog Number: GenBank Accession Number: 10377-1-AP BC001851 GeneID (NCBI): Size: 450 ug/ml 5910 **UNIPROT ID:** Source: Rabbit P52306 Full Name:

RAP1, GTP-GDP dissociation

Immunogen Catalog Number: AG0230 Calculated MW: 66 kDa

> Observed MW: 66 kDa

stimulator 1

Purification Method: Antigen affinity purification Recommended Dilutions:

WB 1:500-1:3000

IHC 1:50-1:500

IP 0.5-4.0 ug for 1.0-3.0 mg of total protein lysate

Applications

Tested Applications: WB, IHC, IP, ELISA **Cited Applications:** WB, IF, IP

Isotype:

Species Specificity: human, mouse, rat **Cited Species:**

human, mouse, chicken

Note-IHC: suggested antigen retrieval with TE buffer pH 9.0; (*) Alternatively, antigen retrieval may be performed with citrate buffer pH 6.0

Positive Controls:

WB: 4T1 cells, HeLa cells, mouse brain tissue, rat brain

tissue

IP: mouse brain tissue, IHC: human gliomas tissue,

Background Information

GDS stimulates the GDP/GTP exchange reaction of a group of small GTP-binding proteins (G proteins), including at least c-Ki-ras p21, smg p21A, smg p21B, rhoA p21 and rhoB p21, by stimulating the dissociation of GDP from and the subsequent binding of GTP to each small G protein. The small GTPase Rap1, which is activated by a large variety of stimuli, functions in the control of integrin-mediated cell adhesion. RAP1GDS1 (synonym: GDS1) is one of RAP1 GTP-GDP dissociation stimulators.

Notable Publications

Author	Pubmed ID	Journal	Application
Jin-Xiao Liang	32968447	Oncol Lett	WB
Beibei Zhang	35138149	Microbiol Spectr	WB,IF
Sahar Nissim	31406347	Nat Genet	IP

Storage


Store at -20°C. Stable for one year after shipment.

Storage Buffer:


PBS with 0.02% sodium azide and 50% glycerol pH 7.3.

Aliquoting is unnecessary for -20°C storage


Selected Validation Data




Various lysates were subjected to SDS PAGE followed by western blot with 10377-1-AP (RAP1GDS1 antibody) at dilution of 1:1500 incubated at room temperature for 1.5 hours.



IP result of anti-RAP1GDS1 (IP:10377-1-AP, 3ug; Detection:10377-1-AP 1:500) with mouse brain tissue lysate 4000ug.



Immunohistochemical analysis of paraffinembedded human gliomas tissue slide using 10377-1-AP (RAP1GDS1 Antibody) at dilution of 1:200 (under 10x lens). Heat mediated antigen retrieval with Tris-EDTA buffer (pH 9.0).



Immunohistochemical analysis of paraffinembedded human gliomas tissue slide using 10377-1-AP (RAP1GDS1 Antibody) at dilution of 1:200 (under 40x lens). Heat mediated antigen retrieval with Tris-EDTA buffer (pH 9.0).