

For Research Use Only

Caspase 9/P35/P10 Polyclonal antibody

Catalog Number: 10380-1-AP

Featured Product

818 Publications



Basic Information

Catalog Number:

10380-1-AP

Concentration:

500 ug/ml

Source:

Rabbit

Isotype:

IgG

Immunogen Catalog Number:

AG0404

GenBank Accession Number:

BC002452

GeneID (NCBI):

842

UNIPROT ID:

P55211

Full Name:

caspase 9, apoptosis-related cysteine peptidase

Calculated MW:

46 kDa

Observed MW:

46 kDa, 30-39 kDa

Purification Method:

Antigen affinity purification

Recommended Dilutions:

WB 1:300-1:1000

IP 0.5-4.0 ug for 1.0-3.0 mg of total protein lysate

IHC 1:50-1:500

IF/ICC 1:200-1:800

Applications

Tested Applications:

WB, IHC, IF/ICC, IP, ELISA

Cited Applications:

WB, IHC, IF, RIP, ELISA

Species Specificity:

human, mouse

Cited Species:

human, mouse, pig, rabbit, canine, chicken, bovine, hamster, goat, duck

Note-IHC: suggested antigen retrieval with TE buffer pH 9.0; (*) Alternatively, antigen retrieval may be performed with citrate buffer pH 6.0

Positive Controls:

WB: Jurkat cells, Staurosporine treated Jurkat cells

IP: HeLa cells,

IHC: mouse lung tissue, human heart tissue

IF/ICC: HepG2 cells,

Background Information

Caspase 9, apoptosis-related cysteine protease (CASP9, synonyms: MCH6, APAF3, APAF-3, ICE-LAP6, CASPASE-9c) is a member of the cysteine-aspartic acid protease (caspase) family. Sequential activation of caspases plays a central role in the execution-phase of cell apoptosis. Caspases exist as inactive proenzymes which undergo proteolytic processing at conserved aspartic residues to produce 2 subunits, large and small, that dimerize to form the active enzyme. Caspase 9 is processed by APAF1; this step is thought to be one of the earliest in the caspase activation cascade. 10380-1-AP can recognize the pre- and cleaved- caspase 9. In recent years, the localization of caspase 9 was a focus of interest. Beside its cytoplasmic distribution, a very extensive localization study was done on rat brain tissue, where caspase 9 was found located predominantly in the nucleus and to a lesser extend in the cytoplasm [PMID: 15541731].

Notable Publications

Author	Pubmed ID	Journal	Application
Xiao-Feng Zhu	36180975	Phytother Res	WB
Faisal Aziz	26427350	Toxicol In Vitro	WB
Zilu Zhang	34570444	Cancer Biol Med	WB

Storage

Storage:

Store at -20°C. Stable for one year after shipment.

Storage Buffer:

PBS with 0.02% sodium azide and 50% glycerol, pH7.3

Aliquoting is unnecessary for -20°C storage

For technical support and original validation data for this product please contact:


T: 4006900926

E: Proteintech-CN@ptglab.com


W: ptgcn.com

This product is exclusively available under Proteintech Group brand and is not available to purchase from any other manufacturer.


Selected Validation Data




Untreated and Staurosporine treated Jurkat cells were subjected to SDS PAGE followed by western blot with 10380-1-AP (Caspase 9/P35/P10 antibody) at dilution of 1:300 incubated at room temperature for 1.5 hours.




IP result of anti-Caspase 9/P35/P10 (IP:10380-1-AP, 3ug; Detection:10380-1-AP 1:200) with HeLa cells lysate 2500 ug.




Immunohistochemical analysis of paraffin-embedded human heart tissue slide using 10380-1-AP (Caspase 9/P35/P10 antibody) at dilution of 1:50 (under 10x lens).




Immunohistochemical analysis of paraffin-embedded human heart tissue slide using 10380-1-AP (Caspase 9/P35/P10 antibody) at dilution of 1:50 (under 40x lens).



Immunohistochemical analysis of paraffin-embedded mouse lung tissue slide using 10380-1-AP (Caspase 9/P35/P10 antibody) at dilution of 1:200 (under 10x lens). Heat mediated antigen retrieval with Tris-EDTA buffer (pH 9.0).



Immunohistochemical analysis of paraffin-embedded mouse lung tissue slide using 10380-1-AP (Caspase 9/P35/P10 antibody) at dilution of 1:200 (under 40x lens). Heat mediated antigen retrieval with Tris-EDTA buffer (pH 9.0).



Immunofluorescent analysis of (-20°C Ethanol) fixed HepG2 cells using Caspase 9/P35/P10 antibody (10380-1-AP) at dilution of 1:400 and Multi-rAb CoraLite @ Plus 488-Goat Anti-Rabbit Recombinant Secondary Antibody (H+L) (RGAR002).