

For Research Use Only

HGS Polyclonal antibody

Catalog Number:10390-1-AP

Featured Product

12 Publications



Basic Information

Catalog Number:

10390-1-AP

Size:

1000 ug/ml

Source:

Rabbit

Isotype:

IgG

Immunogen Catalog Number:

AG0589

GenBank Accession Number:

BC003565

GeneID (NCBI):

9146

UNIPROT ID:

O14964

Full Name:

hepatocyte growth factor-regulated tyrosine kinase substrate

Calculated MW:

86 kDa

Observed MW:

110 kDa

Purification Method:

Antigen affinity purification

Recommended Dilutions:

WB 1:5000-1:50000

IP 0.5-4.0 ug for 1.0-3.0 mg of total protein lysate

IHC 1:50-1:500

IF/ICC 1:200-1:800

Applications

Tested Applications:

WB, IHC, IF/ICC, IP, ELISA

Cited Applications:

WB, IHC, IF

Species Specificity:

human, mouse, rat

Cited Species:

human, mouse

Note-IHC: suggested antigen retrieval with TE buffer pH 9.0; (*) Alternatively, antigen retrieval may be performed with citrate buffer pH 6.0

Positive Controls:

WB : HeLa cells, K-562 cells, mouse brain tissue, rat brain tissue

IP : mouse brain tissue,

IHC : human liver tissue, mouse brain tissue

IF/ICC : HeLa cells,

Background Information

Hepatocyte growth factor-regulated tyrosine kinase substrate (HGS, synonyms: HRS, ZFYVE8) is a 110 to 115-kDa zinc finger phosphotyrosine protein inducible by stimulation with interleukin 2 (IL-2), granulocyte-macrophage colony-stimulating factor (GM-CSF) as well as hepatocyte growth factor (HGF), and is associated with signal-transducing adaptor molecule (STAM). HGS suppresses DNA synthesis upon stimulation with IL-2 and GM-CSF, counteracting STAM's function, which is critical for cell growth signaling mediated by the cytokines. HGS also interacts with the neurofibromatosis 2 tumor suppressor protein Schwannomin/merlin. The growth suppression activity of schwannoma/merlin requires HGS. The binding of schwannoma/merlin to HGS facilitates its ability to function as a tumor suppressor, probably by inhibiting STAT activation.

Notable Publications

Author	Pubmed ID	Journal	Application
Denghui Wei	32958903	Cell Res	WB
Yue-Ming Ling	29042578	Sci Rep	IHC
Jalal M Kazan	34761192	iScience	WB

Storage

Storage:

Store at -20°C. Stable for one year after shipment.

Storage Buffer:

PBS with 0.02% sodium azide and 50% glycerol pH 7.3.

Aliquoting is unnecessary for -20°C storage

For technical support and original validation data for this product please contact:


T: 4006900926

E: Proteintech-CN@ptglab.com


W: ptgcn.com

This product is exclusively available under Proteintech Group brand and is not available to purchase from any other manufacturer.


Selected Validation Data




Various lysates were subjected to SDS PAGE followed by western blot with 10390-1-AP (HGS antibody) at dilution of 1:10000 incubated at room temperature for 1.5 hours.




IP result of anti-HGS (IP:10390-1-AP, 5ug; Detection:10390-1-AP 1:1000) with mouse brain tissue lysate 3000ug.



Immunohistochemical analysis of paraffin-embedded human liver tissue slide using 10390-1-AP (HGS Antibody) at dilution of 1:200 (under 10x lens).



Immunohistochemical analysis of paraffin-embedded human liver tissue slide using 10390-1-AP (HGS Antibody) at dilution of 1:200 (under 40x lens).



Immunofluorescent analysis of (-20°C Ethanol) fixed HeLa cells using HGS antibody (10390-1-AP) at dilution of 1:400 and CoraLite®488-Conjugated AffiniPure Goat Anti-Rabbit IgG(H+L), CL594-Phalloidin (red).