

For Research Use Only

USP2 Polyclonal antibody

Catalog Number: 10392-1-AP

Featured Product

14 Publications



Basic Information

Catalog Number:

10392-1-AP

Size:

400 µg/ml

Source:

Rabbit

Isotype:

IgG

Immunogen Catalog Number:

AG0591

GenBank Accession Number:

BC002955

GeneID (NCBI):

9099

UNIPROT ID:

O75604

Full Name:

ubiquitin specific peptidase 2

Calculated MW:

41 kDa, 69 kDa

Observed MW:

68 kDa, 41-45 kDa

Purification Method:

Antigen affinity purification

Recommended Dilutions:

WB 1:500-1:2000

IHC 1:50-1:500

IF 1:50-1:500

Applications

Tested Applications:

IF/ICC, IHC, WB, ELISA

Cited Applications:

WB, IF, IHC, CoIP

Species Specificity:

human, mouse, rat

Cited Species:

human, rat, mouse

Note-IHC: suggested antigen retrieval with TE buffer pH 9.0; (*) Alternatively, antigen retrieval may be performed with citrate buffer pH 6.0

Positive Controls:

WB: NIH/3T3 cells, mouse brain tissue, mouse kidney tissue, MCF-7 cells, HEK-293 cells

IHC: human breast cancer tissue, human prostate cancer tissue, human skeletal muscle tissue

IF: mouse embryo tissue,

Background Information

Ubiquitin, a highly conserved protein involved in the regulation of intracellular protein breakdown, cell cycle regulation, and stress response, is released from degraded proteins by disassembly of the polyubiquitin chains. The disassembly process is mediated by ubiquitin-specific proteases (USPs). USP2 is also named as UBP41. USP2 can interact indirectly with multiple clock proteins, so it remains possible that USP2 may modulate the ubiquitination and function of these other proteins. It has 4 isoforms produced by alternative splicing. This antibody is specific to the isoform 1(USP2a) of USP2.

Notable Publications

Author	Pubmed ID	Journal	Application
Junhui Xing	32963492	Mediators Inflamm	WB,IHC
Zelong Han	36333288	Cell Death Discov	WB
Jiabei He	30918246	Cell Death Dis	IHC

Storage

Storage:

Store at -20°C. Stable for one year after shipment.

Storage Buffer:

PBS with 0.02% sodium azide and 50% glycerol pH 7.3.

Aliquoting is unnecessary for -20°C storage

For technical support and original validation data for this product please contact:


T: 4006900926

E: Proteintech-CN@ptglab.com


W: ptgcn.com

This product is exclusively available under Proteintech Group brand and is not available to purchase from any other manufacturer.


Selected Validation Data




NIH/3T3 lysates were subjected to SDS PAGE followed by western blot with 10392-1-AP (USP2 antibody) at dilution of 1:1000 incubated at room temperature for 1.5 hours.




Various lysates were subjected to SDS PAGE followed by western blot with 10392-1-AP (USP2 antibody) at dilution of 1:1000 incubated at room temperature for 1.5 hours.




Immunofluorescent analysis of (4% PFA) fixed mouse embryo tissue using 10392-1-AP (USP2 antibody) at dilution of 1:50 and Alexa Fluor 488-conjugated AffiniPure Goat Anti-Rabbit IgG(H+L).



Immunofluorescent analysis of (4% PFA) fixed mouse embryo tissue using 10392-1-AP (USP2 antibody) at dilution of 1:50 and Alexa Fluor 488-conjugated AffiniPure Goat Anti-Rabbit IgG(H+L).



Immunohistochemical analysis of paraffin-embedded human breast cancer tissue slide using 10392-1-AP (USP2 antibody) at dilution of 1:200 (under 10x lens. Heat mediated antigen retrieval with Tris-EDTA buffer (pH 9.0).



Immunohistochemical analysis of paraffin-embedded human breast cancer tissue slide using 10392-1-AP (USP2 antibody) at dilution of 1:200 (under 40x lens. Heat mediated antigen retrieval with Tris-EDTA buffer (pH 9.0).