For Research Use Only

BAP1 Polyclonal antibody

Catalog Number: 10398-1-AP

3 Publications



Basic Information

 Catalog Number:
 GenBank Accession Number:

 10398-1-AP
 BC001596

 Size:
 GeneID (NCBI):

 280 μg/ml
 8314

 Source:
 UNIPROT ID:

 Rabbit
 Q92560

UNIPROT ID: IP 0.5-4.0 ug for 1.0-3.0 mg of total
Q92560 protein lysate
IHC 1:50-1:500

Full Name:
BRCA1 associated protein-1 (ubiquitin

gG

Isotype:

Immunogen Catalog Number:

AG0597

81-91 kDa Observed MW: 91 kDa

Calculated MW:

carboxy-terminal hydrolase)

Applications

Tested Applications: IHC, IP, WB, ELISA Cited Applications: WB

Species Specificity: human, mouse

Cited Species: human, mouse

Note-IHC: suggested antigen retrieval with TE buffer pH 9.0; (*) Alternatively, antigen retrieval may be performed with citrate buffer pH 6.0 Positive Controls:

WB: A2780 cells, human placenta tissue, mouse ovary

Purification Method:

WB 1:500-1:1000

Antigen affinity purification

Recommended Dilutions:

tissue

IP: HeLa cells,

IHC: human ovary tumor tissue, human lung cancer

tissue

Background Information

BRCA1-associated protein-1 (BAP1), also named as KIAA0272, hucep-6, DKFZp686N04275, FLJ35406, FLJ37180, HUCEP-13, is a deubiquitinating enzyme whose function in the control of the cell cycle requires both its deubiquitinating activity and nuclear localization(PMID:22374320). It helps to control cell proliferation by regulating HCFC1 protein levels and by associating with genes involved in the G(1)-S transition)(PMID:19188440). Defects in BAP1 may be a cause of mesothelioma malignant (MESOM)(PMID:21642991) and tumor predisposition syndrome (TPDS)(PMID:21874003). The all BAP1s except those of C. quinquefasciatus have a molecular mass of more than 90 kDa(PMID:22374320).

Notable Publications

Author	Pubmed ID	Journal	Application
Chenggang Zhang	36590171	iScience	WB
Xing Huang	31399419	Cancer Immunol Res	WB
Weiran Ge	36991118	Nature	WB

Storage

Storage:

Store at -20°C. Stable for one year after shipment.

Storage Buffe

PBS with 0.02% sodium azide and 50% glycerol pH 7.3.

Aliquoting is unnecessary for -20°C storage

For technical support and original validation data for this product please contact:

T: 4006900926 E: Proteintech-CN@ptglab.com


W: ptgcn.co

This product is exclusively available under Proteintech Group brand and is not available to purchase from any other manufacturer.


Selected Validation Data



A2780 cells were subjected to SDS PAGE followed by western blot with 10398-1-AP (BAP1 antibody) at dilution of 1:300 incubated at room temperature for 1.5 hours.



Immunohistochemical analysis of paraffinembedded human ovary tumor tissue slide using 10398-1-AP (BAP1 antibody) at dilution of 1:200 (under 10x lens). Heat mediated antigen retrieval with Tris-EDTA buffer (pH 9.0).



Immunohistochemical analysis of paraffinembedded human ovary tumor tissue slide using 10398-1-AP (BAP1 antibody) at dilution of 1:200 (under 40x lens). Heat mediated antigen retrieval with Tris-EDTA buffer (pH 9.0).

IP result of anti-BAP1 (IP:10398-1-AP, 4ug; Detection:10398-1-AP 1:300) with HeLa cells lysate 3000ug.