For Research Use Only

RPA2 Polyclonal antibody

Catalog Number: 10412-1-AP

19 Publications



Basic Information

 Catalog Number:
 GenBank Accession Number:

 10412-1-AP
 BC001630

 Size:
 GeneID (NCBI):

 600 μ g/ml
 6118

 Source:
 UNIPROT ID:

 Rabbit
 P15927

gG replication protein A2, 32kDa

Immunogen Catalog Number:Calculated MW:AG069029 kDa

Observed MW: 32 kDa

Full Name:

Purification Method: Antigen affinity purification

Recommended Dilutions: WB 1:1000-1:6000

IP 0.5-4.0 ug for 1.0-3.0 mg of total

protein lysate IHC 1:20-1:200 IF/ICC 1:50-1:500

Applications

Tested Applications: WB, IHC, IF/ICC, IP, ELISA Cited Applications:

Isotype:

WB, IHC, IF, IP, ChIP Species Specificity: human, mouse, rat Cited Species: human, mouse

Note-IHC: suggested antigen retrieval with TE buffer pH 9.0; (*) Alternatively, antigen retrieval may be performed with citrate buffer pH 6.0

Positive Controls:

WB: HEK-293 cells, mouse brain tissue, HeLa cells, SH-

SY5Y cells

IP: mouse kidney tissue,
IHC: human gliomas tissue,

IF/ICC: A431 cells,

Background Information

RPA2, a subunit of the heterotrimeric replication protein A (RPA) complex, is overexpressed in various cancers. RPA2 is a major target for upstream kinases, including CDKs and the three PIKKs (ATR, ataxia-telangiectasia mutated [ATM], and DNA-PK). RPA2 comprises multiple critical Ser/Tr residues that are phosphorylated in sequential order in response to genotoxic stress.

Notable Publications

Author	Pubmed ID	Journal	Application
Addior	Publiled ID	Journal	Application
Tao Zhang	36050397	Nat Commun	WB
			14/5
Jiao Zhao	35597237	Mol Cell	WB
Yushan Li	35341968	J Genet Genomics	IF
	333,		

Storage

Storage


Store at -20°C. Stable for one year after shipment.

Storage Buffer:


PBS with 0.02% sodium azide and 50% glycerol pH 7.3.

Aliquoting is unnecessary for -20°C storage


Selected Validation Data



Various lysates were subjected to SDS PAGE followed by western blot with 10412-1-Ap (RPA2 antibody) at dilution of 1:3000 incubated at room temperature for 1.5 hours.




IP result of anti-RPA2 (IP:10412-1-Ap, 4ug; Detection:10412-1-Ap 1:1000) with mouse kidney tissue lysate 4000ug.



Immunohistochemical analysis of paraffinembedded human gliomas tissue slide using 10412-1-Ap (RPA2 antibody) at dilution of 1:100 (under 40x lens)..



Immunohistochemical analysis of paraffinembedded human gliomas tissue slide using 10412-1-Ap (RPA2 antibody) at dilution of 1:100 (under 10x lens)..



Immunofluorescent analysis of (4% PFA) fixed A431 cells using RPA2 antibody (10412-1-Ap) at dilution of 1:200 and CoraLite®488-Conjugated AffiniPure Goat Anti-Rabbit IgG(H+L), CL594-phalloidin (red).