

For Research Use Only

# ENOX2 Polyclonal antibody

Catalog Number: 10423-1-AP

Featured Product

11 Publications



## Basic Information

### Catalog Number:

10423-1-AP

### Size:

600 µg/ml

### Source:

Rabbit

### Isotype:

IgG

### Immunogen Catalog Number:

AG0674

### GenBank Accession Number:

BC019254

### GeneID (NCBI):

10495

### UNIPROT ID:

Q16206

### Full Name:

ecto-NOX disulfide-thiol exchanger 2

### Calculated MW:

70 kDa

### Observed MW:

70 kDa

### Purification Method:

Antigen affinity purification

### Recommended Dilutions:

WB 1:500-1:2000

IP 0.5-4.0 µg for 1.0-3.0 mg of total protein lysate

IHC 1:20-1:200

IF/ICC 1:200-1:800

## Applications

### Tested Applications:

WB, IHC, IF/ICC, IP, ELISA

### Cited Applications:

WB, IHC, IP

### Species Specificity:

human, mouse

### Cited Species:

human, mouse

**Note-IHC: suggested antigen retrieval with TE buffer pH 9.0; (\*) Alternatively, antigen retrieval may be performed with citrate buffer pH 6.0**

### Positive Controls:

**WB:** MCF-7 cells, HeLa cells, mouse ovary tissue, PC-3 cells

**IP:** MCF-7 cells,

**IHC:** human ovary tumor tissue,

**IF/ICC:** MCF-7 cells,

## Background Information

Tumor-associated NADH oxidase (ENOX2) as a member of a family of growth-related NADH (or hydroquinone) oxidases. It functions as a terminal oxidase of plasma electron transport from cytosolic NAD(P)H via hydroquinones to acceptors at the cell surface. Hydroquinone oxidase activity alternates with a protein disulfide-thiol interchange/oxidoreductase activity which may control physical membrane displacements associated with vesicle budding or cell enlargement. The activities oscillate with a period length of 22 minutes and play a role in control of the ultradian cellular biological clock

## Notable Publications

Author	Pubmed ID	Journal	Application
Jeng Shiun Chang	36230644	Cancers (Basel)	WB
Atikul Islam	34659883	Am J Cancer Res	IP
Chin-Fang Chang	33163267	Am J Cancer Res	IP

## Storage

### Storage:

Store at -20°C. Stable for one year after shipment.

### Storage Buffer:

PBS with 0.02% sodium azide and 50% glycerol pH 7.3.

Aliquoting is unnecessary for -20°C storage

For technical support and original validation data for this product please contact:


T: 4006900926

E: [Proteintech-CN@ptglab.com](mailto:Proteintech-CN@ptglab.com)


W: [ptgcn.com](http://ptgcn.com)

**This product is exclusively available under Proteintech Group brand and is not available to purchase from any other manufacturer.**


Selected Validation Data




MCF-7 cells were subjected to SDS PAGE followed by western blot with 10423-1-AP (ENOX2 Antibody) at dilution of 1:1000 incubated at room temperature for 1.5 hours.



Immunohistochemical analysis of paraffin-embedded human ovary tumor using 10423-1-AP (ENOX2 antibody) at dilution of 1:100 (under 40x lens).



IP result of anti-ENOX2 (IP:10423-1-AP, 3ug; Detection:10423-1-AP 1:700) with MCF-7 cells lysate 4000ug.



Immunofluorescent analysis of (-20°C Methanol) fixed MCF-7 cells using ENOX2 antibody (10423-1-AP) at dilution of 1:400 and CoraLite®488-Conjugated Goat Anti-Rabbit IgG(H+L) (SA00013-2).