For Research Use Only

Histone H2A type 3 Polyclonal antibody



Catalog Number: 10445-1-AP

Featured Product

4 Publications

Basic Information

Catalog Number: 10445-1-AP Size: 350 μg/ml Source: Rabbit

Isotype: Immunogen Catalog Number:

AG0374

GenBank Accession Number:

BC001193 GeneID (NCBI): 92815 **UNIPROT ID:** Q7L7L0 Full Name:

Calculated MW: 14 kDa Observed MW:

histone cluster 3, H2a

14-18 kDa

Purification Method:

WB 1:500-1:1000

Antigen affinity purification Recommended Dilutions:

Positive Controls:

WB: human brain tissue, fetal human brain tissue

Applications

Tested Applications:

WB, ELISA

Cited Applications: WB, IP, IF, IHC Species Specificity: human, mouse, rat Cited Species: human, mouse

Background Information

Histone H2A type 3, it is expected to be located in the nucleus, and the protein is enriched in the brain tissue. Histones are basic nuclear proteins that are responsible for the nucleosome structure of the chromosomal fiber in eukaryotes. Nucleosomes consist of approximately 146 bp of DNA wrapped around a histone octamer composed of pairs of each of the four core histones (H2A, H2B, H3, and H4). The chromatin fiber is further compacted through the interaction of a linker histone, H1, with the DNA between the nucleosomes to form higher order chromatin structures. This gene is intronless and encodes a replication-dependent histone that is a member of the histone H2A family. The molecular weight of Histone H2A type 3 is 14 kDa.

Notable Publications

Author	Pubmed ID	Journal	Application
Hongyan Ma	34046991	EMBO Rep	IHC
Libo Su	29982651	Nucleic Acids Res	WB,IP,IF
Lihong Yang	38684944	BMC Cancer	WB

Storage

Store at -20°C. Stable for one year after shipment.


PBS with 0.02% sodium azide and 50% glycerol pH 7.3.

Aliquoting is unnecessary for -20°C storage

Selected Validation Data



human brain tissue were subjected to SDS PAGE followed by western blot with 10445-1-AP (Histone H2A type 3 antibody) at dilution of 1:100 incubated at room temperature for 1.5 hours.



fetal human brain tissue were subjected to SDS PAGE followed by western blot with 10445-1-AP (Histone H2A type 3 antibody) at dilution of 1:600 incubated at room temperature for 1.5 hours.