

For Research Use Only

MCM2 Polyclonal antibody

Catalog Number: 10513-1-AP

Featured Product

33 Publications



Basic Information

Catalog Number:

10513-1-AP

Size:

500 µg/ml

Source:

Rabbit

Isotype:

IgG

Immunogen Catalog Number:

AG0798

GenBank Accession Number:

BC007670

GeneID (NCBI):

4171

UNIPROT ID:

P49736

Full Name:

minichromosome maintenance complex component 2

Calculated MW:

102 kDa

Observed MW:

116-125 kDa

Purification Method:

Antigen affinity purification

Recommended Dilutions:

WB 1:500-1:3000

IP 0.5-4.0 µg for 1.0-3.0 mg of total protein lysate

IHC 1:500-1:2000

IF 1:10-1:100

Applications

Tested Applications:

FC, IF/ICC, IHC, IP, WB, ELISA

Cited Applications:

WB, IP, IF, RIP, IHC

Species Specificity:

human, mouse, rat

Cited Species:

human, rat, mouse

Positive Controls:

WB: HEK-293 cells, HeLa cells, K-562 cells

IP: HEK-293 cells,

IHC: human cervical cancer tissue, human tonsillitis tissue

IF: HeLa cells, human gliomas tissue

Note-IHC: suggested antigen retrieval with TE buffer pH 9.0; (*) Alternatively, antigen retrieval may be performed with citrate buffer pH 6.0

Background Information

The MCM2-7 complex forms the core of the replicative helicase which acts as the molecular motor that uses ATP binding and hydrolysis to fuel the unwinding of double-stranded DNA at the replication fork in eukaryotes. This complex is the putative replicative helicase essential for 'once per cell cycle' DNA replication initiation and elongation in eukaryotic cells. MCM2, also named CDCL1 and BM28, is a human nuclear protein that plays an important role in 2 crucial steps of the cell cycle, namely, onset of DNA replication and cell division.

Notable Publications

Author	Pubmed ID	Journal	Application
Irina Vazquez-Villasenor	34536321	Eur J Neurosci	IF
Chun Pan	36115647	Toxicology	IHC
Xin Zhou	36243809	Cell Death Discov	WB,IHC,IF

Storage

Storage:

Store at -20°C. Stable for one year after shipment.

Storage Buffer:

PBS with 0.02% sodium azide and 50% glycerol pH 7.3.

Aliquoting is unnecessary for -20°C storage

For technical support and original validation data for this product please contact:


T: 4006900926

E: Proteintech-CN@ptglab.com


W: ptgcn.com

This product is exclusively available under Proteintech Group brand and is not available to purchase from any other manufacturer.


Selected Validation Data




Various cells were subjected to SDS PAGE followed by western blot with 10513-1-AP (MCM2 antibody) at dilution of 1:1500 incubated at room temperature for 1.5 hours.




WB result of MCM2 antibody (10513-1-AP; 1:2000; incubated at room temperature for 1.5 hours) with sh-Control and sh-MCM2 transfected HEK-293 cells.




Immunofluorescent analysis of (4% PFA) fixed HeLa cells using MCM2 antibody (10513-1-AP) at dilution of 1:20 and CoraLite®488-Conjugated AffiniPure Goat Anti-Rabbit IgG(H+L), CL594-Phalloidin (red).




IP result of anti-MCM2 (IP:10513-1-AP, 4ug; Detection:10513-1-AP 1:800) with HEK-293 cells lysate 2800ug.



Immunohistochemical analysis of paraffin-embedded human cervical cancer tissue slide using 10513-1-AP (MCM2 antibody) at dilution of 1:1000 (under 40x lens). Heat mediated antigen retrieval with Tris-EDTA buffer (pH 9.0).



Immunohistochemical analysis of paraffin-embedded human cervical cancer tissue slide using 10513-1-AP (MCM2 antibody) at dilution of 1:1000 (under 10x lens). Heat mediated antigen retrieval with Tris-EDTA buffer (pH 9.0).



1X10⁶ NIH/3T3 cells were intracellularly stained with 0.2 ug Anti-Human MCM2 (10513-1-AP) and CoraLite®488-Conjugated AffiniPure Goat Anti-Rabbit IgG(H+L) at dilution 1:1000 (red), or 0.2 ug Control Antibody. Cells were fixed and permeabilized with Transcription Factor Staining Buffer Kit (PF00011).