

For Research Use Only

APOD Polyclonal antibody

Catalog Number: 10520-1-AP

Featured Product

5 Publications



Basic Information

Catalog Number:

10520-1-AP

Size:

400 µg/ml

Source:

Rabbit

Isotype:

IgG

Immunogen Catalog Number:

AG0812

GenBank Accession Number:

BC007402

GeneID (NCBI):

347

UNIPROT ID:

P05090

Full Name:

apolipoprotein D

Calculated MW:

33 kDa

Observed MW:

21-33 kDa

Purification Method:

Antigen affinity purification

Recommended Dilutions:

WB 1:500-1:1000

IHC 1:50-1:500

Applications

Tested Applications:

WB, IHC, ELISA

Cited Applications:

WB, IHC

Species Specificity:

human

Cited Species:

human

Note-IHC: suggested antigen retrieval with TE buffer pH 9.0; (*) Alternatively, antigen retrieval may be performed with citrate buffer pH 6.0

Positive Controls:

WB : human serum tissue, HepG2 cells, human plasma

IHC : human liver tissue,

Background Information

Apolipoprotein D (ApoD) is a member of the lipocalin superfamily of ligand transporters, and has been implicated in the transport of small hydrophobic molecules. ApoD is also a component of plasma high-density lipoproteins (HDL). Alteration of ApoD expression has been linked to multiple neurological disorders, including Alzheimer's disease.

Notable Publications

Author	Pubmed ID	Journal	Application
Muhammad Khan	36389725	Front Immunol	IHC
Na Li	34963445	Clin Proteomics	WB
H Bea Kuiperij	31872472	Neuropathol Appl Neurobiol	WB,IHC

Storage

Storage:

Store at -20°C. Stable for one year after shipment.

Storage Buffer:

PBS with 0.02% sodium azide and 50% glycerol pH 7.3.

Aliquoting is unnecessary for -20°C storage

For technical support and original validation data for this product please contact:


T: 4006900926

E: Proteintech-CN@ptglab.com


W: ptgcn.com

This product is exclusively available under Proteintech Group brand and is not available to purchase from any other manufacturer.


Selected Validation Data




human serum were subjected to SDS PAGE followed by western blot with 10520-1-AP (APOD antibody) at dilution of 1:1000 incubated at room temperature for 1.5 hours.



HepG2 cells were subjected to SDS PAGE followed by western blot with 10520-1-AP (APOD antibody) at dilution of 1:1000 incubated at room temperature for 1.5 hours.



Immunohistochemical analysis of paraffin-embedded human liver tissue slide using 10520-1-AP (APOD antibody) at dilution of 1:200 (under 10x lens). Heat mediated antigen retrieval with Tris-EDTA buffer (pH 9.0).



Immunohistochemical analysis of paraffin-embedded human liver tissue slide using 10520-1-AP (APOD antibody) at dilution of 1:200 (under 40x lens). Heat mediated antigen retrieval with Tris-EDTA buffer (pH 9.0).