

For Research Use Only

MAOA Polyclonal antibody

Catalog Number: 10539-1-AP

Featured Product

27 Publications



Basic Information

Catalog Number:

10539-1-AP

Concentration:

800 ug/ml

Source:

Rabbit

Isotype:

IgG

Immunogen Catalog Number:

AG0821

GenBank Accession Number:

BC008064

GeneID (NCBI):

4128

UNIPROT ID:

P21397

Full Name:

monoamine oxidase A

Calculated MW:

60 kDa

Observed MW:

60 kDa

Purification Method:

Antigen affinity purification

Recommended Dilutions:

WB 1:500-1:3000

IP 0.5-4.0 ug for 1.0-3.0 mg of total protein lysate

IHC 1:300-1:1200

Applications

Tested Applications:

WB, IHC, IP, ELISA

Cited Applications:

WB, IHC, IF, CoIP

Species Specificity:

human

Cited Species:

human, mouse, rat

Note-IHC: suggested antigen retrieval with TE buffer pH 9.0; (*) Alternatively, antigen retrieval may be performed with citrate buffer pH 6.0

Positive Controls:

WB : LNCaP cells, human placenta tissue

IP : LNCaP cells,

IHC : human pancreas cancer tissue, human kidney tissue, human liver tissue, human pancreas tissue

Background Information

MAOA belongs to the flavin monoamine oxidase family. It catalyzes the oxidative deamination of biogenic and xenobiotic amines and has important functions in the metabolism of neuroactive and vasoactive amines in the central nervous system and peripheral tissues. MAOA preferentially oxidizes biogenic amines such as 5-hydroxytryptamine (5-HT), norepinephrine and epinephrine.

Notable Publications

Author	Pubmed ID	Journal	Application
Xiaochao Tan	32931483	J Clin Invest	
Odetta Antico	34767452	Sci Adv	WB
Yinrui Guo	34045460	Transl Psychiatry	WB

Storage

Storage:

Store at -20°C. Stable for one year after shipment.

Storage Buffer:

PBS with 0.02% sodium azide and 50% glycerol pH 7.3.

Aliquoting is unnecessary for -20°C storage

For technical support and original validation data for this product please contact:


T: 4006900926

E: Proteintech-CN@ptglab.com


W: ptgcn.com

This product is exclusively available under Proteintech Group brand and is not available to purchase from any other manufacturer.


Selected Validation Data




LNCaP cells were subjected to SDS PAGE followed by western blot with 10539-1-AP (MAOA antibody) at dilution of 1:1500 incubated at room temperature for 1.5 hours.




Immunohistochemical analysis of paraffin-embedded human pancreas cancer tissue slide using 10539-1-AP (MAOA antibody) at dilution of 1:600 (under 10x lens). Heat mediated antigen retrieval with Tris-EDTA buffer (pH 9.0).



Immunohistochemical analysis of paraffin-embedded human pancreas cancer tissue slide using 10539-1-AP (MAOA antibody) at dilution of 1:600 (under 40x lens). Heat mediated antigen retrieval with Tris-EDTA buffer (pH 9.0).



IP result of anti-MAOA (IP:10539-1-AP, 4ug; Detection:10539-1-AP 1:5000) with LNCaP cells lysate 880 ug.



Immunohistochemical analysis of paraffin-embedded human pancreas cancer tissue slide using 10539-1-AP (MAOA antibody) at dilution of 1:600 (under 40x lens). Heat mediated antigen retrieval with Tris-EDTA buffer (pH 9.0).