For Research Use Only

GTF2A2 Polyclonal antibody

Catalog Number: 10540-1-AP



Basic Information

Catalog Number:

10540-1-AP

Size:

400 ug/ml

Source:

Rabbit

BC001919

GeneID (NCBI):

2958

Surroe:

UNIPROT ID:

Rabbit

P52657

Isotype:

GeneID (NCBI):

Full Name:

general transcription factor IIA, 2,

Immunogen Catalog Number: 12kDa

AG0825 Calculated MW:

12 kDa
Observed MW:
12 kDa

Applications

Tested Applications: IHC, WB, ELISA
Species Specificity:

IgG

human, mouse, rat

Note-IHC: suggested antigen retrieval with TE buffer pH 9.0; (*) Alternatively, antigen retrieval may be performed with citrate buffer pH 6.0

Background Information

Transcription mediated by RNA polymerase II depends on a set of different transcription factors to form the preinitiation complex. TFIIA is involved in the construction of this complex and increases the affinity of TBP for the DNA union region [PMID:20480367]. TFIIA in a complex with TBP mediates transcriptional activity [PMID: 11030333]. GTF2A2 is one subunit of TFIIA.

Storage

Storage:

Store at -20°C. Stable for one year after shipment.

Storage Buffer:

PBS with 0.02% sodium azide and 50% glycerol pH 7.3.

Aliquoting is unnecessary for -20°C storage

Positive Controls:

WB: mouse testis tissue. HeLa cells

IHC: mouse liver tissue, mouse heart tissue, rat liver

Purification Method:

WB 1:200-1:1000 IHC 1:500-1:2000

Antigen affinity purification


Recommended Dilutions:

tissue

Selected Validation Data



mouse testis tissue were subjected to SDS PAGE followed by western blot with 10540-1-AP (GTF2A2 antibody) at dilution of 1:300 incubated at room temperature for 1.5 hours.



Immunohistochemical analysis of paraffinembedded mouse liver tissue slide using 10540-1-AP (GTF2A2 antibody) at dilution of 1:1000 (under 40x lens). Heat mediated antigen retrieval with Tris-EDTA buffer (pH 9.0).