For Research Use Only

GTF2A2 Polyclonal antibody

Catalog Number: 10540-1-AP



Purification Method:

WB 1:200-1:1000 IHC 1:20-1:200

Antigen affinity purification

Recommended Dilutions:

Basic Information

Catalog Number: GenBank Accession Number: 10540-1-AP BC001919 GeneID (NCBI): Size: 400 µg/ml 2958 **UNIPROT ID:** Source: Rabbit P52657 Full Name: Isotype:

general transcription factor IIA, 2,

12kDa Immunogen Catalog Number:

AG0825 Calculated MW:

12 kDa Observed MW: 12 kDa

Applications

Tested Applications: IHC, WB, ELISA Species Specificity: human, mouse, rat

Note-IHC: suggested antigen retrieval with TE buffer pH 9.0; (*) Alternatively, antigen retrieval may be performed with citrate buffer pH 6.0

Background Information

Transcription mediated by RNA polymerase II depends on a set of different transcription factors to form the preinitiation complex. TFIIA is involved in the construction of this complex and increases the affinity of TBP for the DNA union region [PMID:20480367]. TFIIA in a complex with TBP mediates transcriptional activity [PMID: 11030333]. GTF2A2 is one subunit of TFIIA.

Positive Controls:

WB: mouse testis tissue. HeLa cells

IHC: human prostate cancer tissue,

Storage

Storage:

Store at -20°C. Stable for one year after shipment.

Storage Buffer:


PBS with 0.02% sodium azide and 50% glycerol pH 7.3.

Aliquoting is unnecessary for -20°C storage

Selected Validation Data



mouse testis tissue were subjected to SDS PAGE followed by western blot with 10540-1-AP (GTF2A2 antibody) at dilution of 1:300 incubated at room temperature for 1.5 hours.



Immunohistochemical analysis of paraffinembedded human prostate cancer using 10540-1-AP (GTF2A2 antibody) at dilution of 1:100 (under 25x lens).