

For Research Use Only

OTUB1 Polyclonal antibody, PBS Only

Catalog Number: 10573-1-PBS

Featured Product



Basic Information

Catalog Number:

10573-1-PBS

Concentration:

1 mg/ml

Source:

Rabbit

Isotype:

IgG

Immunogen Catalog Number:

AG0845

GenBank Accession Number:

BC007519

GeneID (NCBI):

55611

UNIPROT ID:

Q96FW1

Full Name:

OTU domain, ubiquitin aldehyde binding 1

Calculated MW:

31 kDa and 35 kDa

Observed MW:

31-35 kDa

Purification Method:

Antigen affinity purification

Applications

Tested Applications:

WB, IF/ICC, Indirect ELISA

Species Specificity:

human, mouse, rat

Background Information

OTUB1 belongs to the ovarian tumor domain protease (OTU) subfamily of deubiquitinases, which could block ubiquitination leading to protein stability. OTUB1 mainly prefers k-48 and k-63 ubiquitin chains type, especially in k-48 ubiquitin chain. OTUB1 performs its deubiquitinating function via two distinct manners: the canonical and non-canonical manner. OTUB1 is involved in many critical immune response processes, containing CD8 T cell activation, PD-L1 related immune evasion, and several infectious diseases. OTUB1 plays an important role in the occurrence, proliferation, invasion, and progression of various cancers, such as bladder cancer, breast cancer, prostate cancer, and colorectal cancer (PMID: 34875341, 35296795). OTUB1 has 2 isoforms with molecular weights of 31 and 35 kDa, respectively. With phospho or ubiquitin modification or reaction, the MW of OTUB1 will be increased.

Storage

Storage:

Store at -80°C.

The product is shipped with ice packs. Upon receipt, store it immediately at -80°C

Storage Buffer:

PBS only, pH7.3

For technical support and original validation data for this product please contact:


T: 4006900926

E: Proteintech-CN@ptglab.com


W: ptgcn.com

This product is exclusively available under Proteintech Group brand and is not available to purchase from any other manufacturer.


Selected Validation Data



Various lysates were subjected to SDS PAGE followed by western blot with 10573-1-AP (OTUB1 antibody) at dilution of 1:1500 incubated at room temperature for 1.5 hours. This data was developed using the same antibody clone with 10573-1-PBS in a different storage buffer formulation.



WB result of OTUB1 antibody (10573-1-AP; 1:10000; incubated at room temperature for 1.5 hours) with sh-Control and sh-OTUB1 transfected HepG2 cells. This data was developed using the same antibody clone with 10573-1-PBS in a different storage buffer formulation.



Immunofluorescent analysis of (-20°C Ethanol) fixed U2OS cells using OTUB1 antibody (10573-1-AP) at dilution of 1:400 and CoraLite® 488-Conjugated AffiniPure Goat Anti-Rabbit IgG(H+L) (SA00013-2), CL594-phalloidin (red). This data was developed using the same antibody clone with 10573-1-PBS in a different storage buffer formulation.