For Research Use Only

T: 4006900926

E: Proteintech-CN@ptglab.com

## ERCC3 Polyclonal antibody Catalog Number: 10580-1-AP Featured Product





Group brand and is not available to purchase from any


other manufacturer.

Basic Information	Catalog Number: 10580-1-AP	GenBank Accession Numl BC008820	per: Purification Me Antigen affinit		
	Size:	GeneID (NCBI):	Recommended		
	350 ug/ml	2071	WB 1:500-1:100		
	Source:	UNIPROT ID:	•	r 1.0-3.0 mg of total	
	Rabbit	P19447	protein lysate IHC 1:50-1:500		
	Isotype:	Full Name:			
	IgG Immunogen Catalog Number: AG0876	excision repair cross-complementing rodent repair deficiency, complementation group 3 (xeroderma pigmentosum group B			
	complementing)				
		Calculated MW: 89 kDa			
		Observed MW: 89 kDa			
Applications	Tested Applications:	Pc	Positive Controls:		
	WB, IHC, IP, ELISA		WB : HEK-293 cells, HeLa cells, A431 cells, COLO 320		
	Cited Applications: WB		ells, HT-1080 cells, mouse co	lon tissue	
	Species Specificity:	IP	IP : A431 cells,		
	human, mouse, rat	IH	IC : human stomach tissue,		
	Cited Species: human				
	Note-IHC: suggested antigen retrieval with TE buffer pH 9.0; (*) Alternatively, antigen retrieval may be performed with citrate buffer pH 6.0				
		l with citrate			
Background Information		ting 3/xeroderma pigmentosu ttions as an integral componer nscription and global genome lular level, a defect in ERCC3	nt of the transcription factor repair as well as in transcrip results in high UV sensitivit	IIH complex. ERCC3 i ption-coupled repair	
	<b>buffer pH 6.0</b> Excision repair cross-complement dependent DNA helicase that func involved in initiation of basal trai (TCR) [PMID:19942596]. At the cell synthesis recovery and decreased	ting 3/xeroderma pigmentosu ttions as an integral componer nscription and global genome lular level, a defect in ERCC3	nt of the transcription factor repair as well as in transcrip results in high UV sensitivit	IIH complex. ERCC3 i ption-coupled repair	
	buffer pH 6.0 Excision repair cross-complement dependent DNA helicase that func involved in initiation of basal trai (TCR) [PMID:19942596]. At the cel synthesis recovery and decreased	ting 3/xeroderma pigmentosu tions as an integral component nscription and global genome lular level, a defect in ERCC3 DNA repair due to defective N Pubmed ID Journal	nt of the transcription factor repair as well as in transcrip results in high UV sensitivit	IIH complex. ERCC3 i btion-coupled repair y, reduced post-UV RN	
Background Information	buffer pH 6.0 Excision repair cross-complement dependent DNA helicase that func involved in initiation of basal trai (TCR) [PMID:19942596]. At the cel synthesis recovery and decreased	ting 3/xeroderma pigmentosu tions as an integral component nscription and global genome lular level, a defect in ERCC3 DNA repair due to defective N Pubmed ID Journal	nt of the transcription factor repair as well as in transcrip results in high UV sensitivit IER [PMID:16947863]. Biophys Res Commun	IIH complex. ERCC3 i otion-coupled repair y, reduced post-UV RM Application	
	buffer pH 6.0 Excision repair cross-complement dependent DNA helicase that func involved in initiation of basal trai (TCR) [PMID:19942596]. At the cell synthesis recovery and decreased Author Xie Qiong-Hui QH	ting 3/xeroderma pigmentosu ttions as an integral componen nscription and global genome lular level, a defect in ERCC3 DNA repair due to defective N Pubmed ID Journal 21672525 Biochem	nt of the transcription factor repair as well as in transcrip results in high UV sensitivit IER [PMID:16947863]. Biophys Res Commun	IIH complex. ERCC3 i otion-coupled repair y, reduced post-UV RM Application WB	

W: ptgcn.com

## Selected Validation Data





Immunohistochemical analysis of paraffinembedded human stomach tissue slide using 10580-1-AP (ERCC3 antibody) at dilution of 1:200 (under 20x lens). Heat mediated antigen retrieval with Tris-EDTA buffer (pH 9.0).

HEK-293 cells were subjected to SDS PAGE followed by western blot with 10580-1-AP (ERCC3 antibody) at dilution of 1:500 incubated at room temperature for 1.5 hours. IP result of anti-ERCC3 (IP:10580-1-AP, 4ug; Detection:10580-1-AP 1:500) with A431 cells lysate 1200ug.